

ANSWER KEY & SOLUTION KEY FINAL ROUND - 02 (PCB) Dt.03.04.2024

BOTANY

SECTION - A (35 Questions)

01. (3) (11th, para 8.4.1, 8.4.2, page no. 128,129)
02. (4) (NCERT XI added Family by NMC)
03. (3) (NCERT XI Pg.230, 14.3)
04. (4) [NCERT XI Pg. No. 222, 13.10]
05. (2) [NCERT class XI, Page no. 93, Point no. 6.3.4]
06. (1) (NCERT XI Pg.236, 14.7)
07. (3) [NCERT XI Pg. No. 216, 13.7.1 and page no. 219 2nd paragraph]
08. (3) (NCERT XII, Pg 80, 5.3.1, based on Law of Independent Assortment)
09. (4) (NCERT XII, Pg 101, 6.2.1)
10. (2) (NCERT 12th, Page No-29, Paragraph- 2, Line No- 7 to 10)
11. (4) (NCERT 12th, Page No-21, Paragraph- 2, Line No- 9 to 11)
12. (3) (NCERT XII, Pg 89- 90. Based on sex linkage)
13. (4) (NCERT page no.38 gymno.concept)
14. (2) (11th NCERT Page no.28 to 29, conceptual)
15. (2) (NCERT page no. 34 ,3.2 concept)
16. (1) (11th, para10.4 ,page no. 167)
17. (4) [NCERT XI, Page 248, Point 15.4.3.1]
18. (4) (NCERT XI Page No. 77, sub-topic 5.7.2)
19. (4) (12NCERT,Page no.246 fig.14.2)
20. (4) (12thNCERT old page no.243 3rd para)
21. (2) (NCERT 11th, Page No-11, Table-1.1)
22. (3) (NCERT XII, Pg 75, Para 1, Line 4)
23. (4) (NCERT XII, Pg 78, Dihybrid cross based)
24. (1) (11th, para 10.1.1, Figure 10.1, page no. 163)
25. (4) (NCERT 11th, Page No-18, Figure 2.1)
26. (3) (NCERT 11th, Page No- 22, Paragraph-2, Line No- 4,5)
27. (1) (NCERT 11th, Page No-19, Paragraph- 2.1.2, Line No-27,28)
28. (2) [NCERT Page no. 87 (Point 6.1.2.2), 88 (Line number- 01-20), 90 (Line no. 01-03)]
29. (1) (NCERT 12th, Page No-35, Paragraph- 2.4.2, Line No- 13-15)
30. (3) (NCERT XI Page No. 75, sub-topic 5.5.1.3 & added family)

31. (3) (NCERT XII, Pg118, Para-5 , Line-1)
32. (2) (NCERT XII, Page No. 109, Para 3, Line 5-6)
33. (4) (NCERT XII, Pg 102, Figure 6.5 The Hershey-Chase experiment)
34. (1) [NCERT XI, Page 248 (First paragraph), 249 (Last paragraph)]
35. (3) (11th, para 8.2, page no. 126)

SECTION - B (Attempt Any 10 Questions)

36. (4) (NCERT XII, Pg.101, Para 5, Line 2)
37. (4) (11th, para 8.4.1,8.5.7, 8.5.6, 8.5.9, page no.126, 136,137)
38. (4) (NCERT 12th, Page No-23, Paragraph- 2, Line No- 15-17)
39. (3) (NCERT XII, Pg. 78 , Para 1, Line 4)
40. (3) (NCERT 11th, Page No-22, Last Paragraph, Line No-27,28)
41. (4) (NCERT XII, Pg 111, Para 2)
42. (1) (11th, para 10.2.1, page no. 164) (11th, para 8.5.1, page no. 131)
43. (2) (NCERT XII, Pg 91, 5.8.3)
44. (4) NCERT page no.38 ,1st para, 1st line
45. (2) (NCERT 11th, Page No-9, Paragraph- 1.3.3, Line No-31,32)
46. (4) [NCERT Page no. 89, Second paragraph]
47. (4) (NCERT XI Page No. 80, sub-topic 5.9.2)
48. (2) (NCERT XI Pg.233, Fig.14.4)
49. (4) (11th, para 8.3, page no. 127)
50. (3) [NCERT XI Pg. No.210, 13.4, fig 13.3a based]

ZOOLOGY

SECTION - A (35 Questions)

51. (3) (NCERT XI Page No. 47, 4.1.2)
52. (1) (NCERT XI Page No. 51, 4.2.4)
53. (1) (NCERT XI Page No. 55, table 4.1)
54. (3) (NCERT 11th, Page No- 143, Table- 9.1)
55. (1) (NCERT 11th, Page No- 151, 1st Paragraph, Line No-1 to 12)
56. (2) (NCERT 11th page 119, para3, line)
57. (3) (NCERT Pg.No.284- Cardiac cycle)
58. (1) (NCERT XI Page No. 294, 6th paragraph)

59. (4) (NCERT 12th, p.no. 52, para 1)
 60. (2) (NCERT 12th, p.no. 53, para 1, line 1)
 61. (1) (NCERT 12th, p.no. 43, para 3)
 62. (1) (NCERT 12th p.no. 59)
 63. (3) (NCERT 11th page no 112, para 1, line 6)
 64. (3) (NCERT Pg.No.153 - Immune system)
 65. (4) (NCERT Pg.No.156 - Cancer)
 66. (3) (NCERT Pg. No.-146)
 67. (3) [NCERT P. No.309 3rd line below Diagram]
 68. (2) [NCERT P. No.321 1st para last 3 lines]
 69. (1) [NCERT P. No.318 Para Below Diagram]
 70. (4) [NCERT P. No.212 3rd para]
 71. (1) (NCERT Pg.No.-274)
 72. (1) (NCERT Pg. No.-161)
 73. (2) [NCERT P. No.311 Last line of Pelvic Girdle para]
 74. (1) [NCERT P. No. 213- 3rd Point, P No 201-2nd para , 11.3 & 11.3.1,]
 75. (4) [NCERT P. No. 213- 2nd, 3rd & 4th Point]
 76. (1) [NCERT P. No.311 1st Para]
 77. (2) (12th, para 10.4 , page no.185/zoology.)
 78. (1) (12th, para10.2.3 , page no.183/zoology.)
 79. (2) (12th ncert 10.1 BIOTECHNOLOGICAL APPLICATIONS IN AGRICULTURE)
 80. (2) (NCERT Pg.No. 272- Lung Volumes)
 81. (3) (NCERT page no.260, fig.15.1)
 82. (4) (NCERT XI Chapter chemical co-ordination conceptual)
 83. (1) (NCERT 12th, Page No-130, 1st Paragraph, Line No- 7 to 9)
 84. (3) NCERT 12th, Page No-135, 3rd Paragraph, Line No- 5,6
 85. (3) (Old 12th NCERT page no.232 1st para.)

SECTION - B (Attempt Any 10 Questions)

86. (2) [NCERT P. No.198 3rd para]
 87. (4) (NCERT Pg.No.284- Cardiac cycle)
 88. (4) [NCERT Practical Syllabus P. No.125 4th Point and last Para]
 89. (4) (NCERT XI Page No. 293, 2nd paragraph)
 90. (3) (NCERT XI Page No. 340, first paragraph)
 91. (4) (NCERT 12th p.no.64, para 3, LINE 5)
 92. (2) (NCERT 12th p.no. 57, para 2, line 4)
 93. (1) (NCERT 11th, Page No-148, Figure- 9.2, Conceptual)
 94. (4) (NCERT 12th, Page No-127, Last Paragraph, Line No- 1,2 and 3)
 95. (3) (NCERT page no.259; 2nd para)
 96. (4) (12th NCERT, Page no.230 to 231)
 97. (1) [NCERT P. No.317 1st para first 4 lines]
 98. (1) NCERT 12th, Page No-140, 2nd Paragraph, Line No- 5 to 8
 99. (3) (NCERT 11th, page 101, para 3, line 2)
 100. (3) (NCERT Pg. No.158)

PHYSICS


SECTION - A (35 Questions)

101. (1)
 102. (2) $\vec{r} = at^2\hat{i} + bt\hat{j}$... (i)
 $x = at^2$... (ii)
 and $y = bt$
 From Eq. (ii) put value of t in Eq. (i)

$$x = \frac{a}{b^2} y^2$$


 103. (4)
 104. (1) As we know, momentum, $\vec{p} = mv$
 Change in P can be brought by changing force \vec{F} i.e.
 As, $\vec{F} = \frac{dp}{dt}$ rate of change of momentum with time.
 \Rightarrow as $\Delta p = F\Delta t = \text{impulse}$ so for same F and Δt , Δp will be same even for different bodies.
 105. (4) Work done, $W = \int F \cdot dy = \int_0^1 (20 + 10y) dy$

$$= \left[20y + \frac{10}{2} y^2 \right] = 20 + \frac{10}{2} = 25 \text{ J}$$



 106. (2)
 $u = \sqrt{2gh_1}$, $v = \sqrt{2gh_2}$
 $\Delta p = m(v + u)$
 $= m(\sqrt{2gh_2} + \sqrt{2gh_1})$
 107. (3) As $CD = 2AB$, latent heat of vaporization is twice the latent heat of fusion.
 \therefore Statement (3) is false. All other statements are correct.
 108. (4) $X = 3YZ^2$
 $\therefore [Y] = \frac{[X]}{[Z]^2} = \frac{[M^{-1}L^{-2}T^4A^2]}{[MT^{-2}A^{-1}]^2} = [M^{-3}L^{-2}T^8A^4]$
 109. (1) According to Wien's displacement law
 $\lambda_m \propto \frac{1}{T} \Rightarrow \lambda_{m_2} < \lambda_{m_1}$ ($\because T_1 < T_2$)
 Therefore, $I - \lambda$ graph for T_2 have lesser wavelength (λ_m) and so curve for T_2 will shift towards left side.
 110. (3) $U = nC_V T$

$$\begin{aligned} \Rightarrow U &= n_1 C_{V_1} T + n_2 C_{V_2} T \\ &= 8 \times \frac{3R}{2} \times T + 6 \times \frac{5R}{2} \times T \\ &= 27RT. \end{aligned}$$

111. (2) $h = R \cos \theta = 12 \times \frac{2}{3} \text{ cm}$



The equivalent circuit diagrams are as shown in the figure below.



The equivalent capacitance between A and B is

$$C_{AB} = \frac{2C}{3} + \frac{2C}{3} = \frac{4C}{3}$$

113. (2) When cube is of side a and point charge Q is at the center of the cube then the total electric flux due to this charge will pass evenly through the six faces of the cube. So, the electric flux through one face will be equal to 1/6 of the total electric flux due to this charge.

$$\text{Flux through 6 faces} = \frac{Q}{\epsilon_0}$$

$$\therefore \text{Flux through 1 face,} = \frac{Q}{6\epsilon_0}$$

114. (1) $v_d = \frac{e}{m} \times \frac{V}{l} \tau$ or $v_d = \frac{e}{m} \cdot \frac{El}{l} \tau$ (since $V = El$)

$$\therefore v_d \propto E$$

115. (1) $B = \frac{E}{v}$

$$\begin{aligned} &= \frac{600\text{V}}{3 \times 10^{-3} \text{ m}} \times \frac{1}{2 \times 10^6 \text{ ms}^{-1}} \\ &= 10^{-1} \text{ T} = 0.1 \text{ T} \end{aligned}$$

116. (1) Magnetic field inside the hollow metallic cylinder

$$B_{in} = 0, \text{ and magnetic field outside it } B_{out} \propto \frac{1}{r}$$

117. (2) Eddy currents may be reduced by using laminated core of soft iron.

118. (3) $I = \frac{V}{X_C}$ in Pure Capacitor

$$\begin{aligned} &= \frac{V}{\frac{1}{2\pi f C}} = V 2\pi f C \\ &\Rightarrow I \propto f \end{aligned}$$

119. (4) (A) \rightarrow (2); (B) \rightarrow (1); (C) \rightarrow (4); (D) \rightarrow (3)

120. (1)

121. (3)

122. (4) Mass defect = $\frac{B.E.}{c^2}$

Mass of nucleus = Mass of proton + mass of neutron – mass defect.

123. (1)

124. (2)

125. (2) As current is flowing from A to B, therefore, moving from A to B,

$$V_A - V_B = [5 \times 1 - 15 + 5 \times 10^{-3} \times (-10^3)] = -15 \text{ volt}$$

$$\therefore V_B - V_A = 15 \text{ V}$$

126. (1) Wave function should be in the form $y = f(x, t)$

For travelling wave y should be linear function of x and t and they must exist as $(x \pm vt)$.

$$y = A \sin(15x - 2t)$$

127. (4) Speed of light does not depend on the motion of source as well as intensity.

128. (4) For telescope

$$\text{Tube length } (L) = f_o + f_e = 60$$

$$\text{and magnification } (m) = \frac{f_o}{f_e} = 5 \Rightarrow f_o = 5f_e$$

$$\therefore f_o = 50 \text{ cm and } f_e = 10 \text{ cm}$$

Hence focal length of eye-piece, $f_e = 10 \text{ cm}$.

129. (4) Let μ_0 = Mass of body in vacuum

Weight of body in air

= Weight of standard weights in air

$$\therefore \mu_0 g - \left(\frac{\mu_0}{d_1} \right) dg = Mg - \frac{M}{d_2} dg$$

$$\text{or } \mu_0 = \frac{M \left(1 - \frac{d}{d_2} \right)}{\left(1 - \frac{d}{d_1} \right)}$$

Hence, the correct answer is option (4)

130. (2) $\lambda_{med} = \frac{\lambda_{vacuum}}{\mu}$

and we know that fringe width $\beta = \frac{\lambda D}{d}$

Therefore, $\beta_{med} = \frac{\beta_{vacuum}}{\mu} = \frac{12}{4/3} = 9 \text{ mm}$

131. (1)

132. (3) Two particles will meet at their centre of mass
 \therefore Distance of the centre of mass from 8 kg mass

$$= \frac{8 \times 0 + 4 \times 12}{8 + 4} = 4m.$$

133. (2) Zero error is positive because CS division is ahead of pitch scale making.

Least count =

$$\frac{\text{Pitch}}{\text{No. of divisions on circular scale}} = \frac{0.5 \text{ mm}}{50} = 1 \times 10^{-5} \text{ m} = 10 \mu\text{m}.$$


134. (2) When a child sits on a rotating disc, no external torque is introduced. Hence the angular momentum of the system is conserved. But the moment of inertia of the system will increase and as a result, the angular speed of the disc will decrease to maintain constant angular momentum. (2)

135. (2) $[\tau] = [ML^2 T^{-2}]$

$$\frac{\Delta\tau}{\tau} \times 100 = \frac{\Delta M}{M} \times 100 + 2 \frac{\Delta L}{L} \times 100 + 2 \frac{\Delta T}{T} \times 100 = 5\% + 2 \times 5\% + 2 \times 5\% = 25\%$$

SECTION - B (Attempt Any 10 Questions)

136. (2)



For M block

$$10g \sin 53^\circ - \mu (10g) \cos 53^\circ - T = 10 \times 2$$

$$T = 80 - 15 - 20$$

$$T = 45 \text{ N}$$

For m block

$$T - mg \sin 37^\circ - \mu mg \cos 37^\circ = m \times 2$$

$$45 = 10m$$

$$m = 4.5 \text{ kg}$$

137. (3) $f = kx = Mg$

$$\frac{1}{2} kx^2 = mgx$$

$$\Rightarrow kx = 2mg$$

$$\Rightarrow Mg = 2mg \Rightarrow m = \frac{M}{2}$$

138. (4) According to Einstein's photoelectric equation

$$K_{max} = hv - \phi$$

$$eV_s = \frac{hc}{\lambda} - \phi_0 \Rightarrow V_s = \frac{hc}{\lambda e} - \frac{\phi_0}{e}$$

where, λ - Wavelength of incident light

ϕ_0 - Work function

V_s - Stopping potential

According to given problem,

$$V_1 = \frac{hc}{\lambda e} - \frac{\phi_0}{e} \dots\dots(i)$$

$$V_2 = \frac{hc}{(\lambda/2)e} - \frac{\phi_0}{e}$$


$$V_2 = \frac{2hc}{\lambda e} - \frac{\phi_0}{e} = \frac{2hc}{\lambda e} - \frac{2\phi_0}{e} + \frac{2\phi_0}{e} - \frac{\phi_0}{e}$$

$$= 2 \left(\frac{hc}{\lambda e} - \frac{\phi_0}{e} \right) + \frac{\phi_0}{e}$$

$$V_2 = 2V_1 + f_e \text{ [Using eqn. (i)]}$$

$$\therefore V_2 > 2V_1.$$

139. (2) The charge distribution in the surfaces of the concentric spherical shells is shown in the figure.



$$\text{Charge density, } \sigma = \frac{\text{Charge}}{\text{Area}}$$

Since the surface charge densities are equal

$$\therefore \frac{Q_1}{4\pi R^2} = \frac{Q_2}{4\pi(4R^2)} = \frac{Q_3}{4\pi(9R^2)}$$

$$Q_1 = \frac{Q_2 + Q_3}{4} \dots(i)$$

$$\text{and } Q_1 = \frac{Q_1 + Q_2 + Q_3}{9} \dots(ii)$$

$$\text{From (i), we get } 3Q_1 = Q_2 \dots(iii)$$

From (ii) we get

$$8Q_1 - Q_2 = Q_3$$

$$8Q_1 - 3Q_1 = Q_3 \text{ Using (iii)}$$

$$5Q_1 = Q_3$$

$$\therefore \frac{Q_1}{1} = \frac{Q_2}{3} = \frac{Q_3}{5} \text{ or } Q_1 : Q_2 : Q_3 = 1 : 3 : 5$$

140. (3) The gravitational field due to the ring at a distance $\sqrt{3}r$ is given by

$$E = \frac{Gm(\sqrt{3}r)}{[r^2 + (\sqrt{3}r)^2]^{3/2}} \Rightarrow E = \frac{\sqrt{3}Gm}{8r^2}$$

$$\text{Attractive force} = \sqrt{3} \frac{GMm}{8r^2}$$

141. (2) Max. KE of a particle in SHM

$$= \frac{1}{2}kr^2 = \frac{1}{2} \times 200 \times 1^2 = 100\text{J}$$

As total energy = Max. KE + Minimum PE

$$\therefore 150 = 100 + \text{minimum PE}$$

$$\text{or Minimum PE} = 150 - 100 = 50\text{ J}$$

Thus, both, Statement I and Statement II are false.

142. (4) Let h be the height of liquid surface in the vessel. The velocity of efflux is given by

$$v_{\text{eff}} = \sqrt{2gh}$$

If H be the height of table, then

$$H = \frac{1}{2}gt^2 \text{ or } t = \sqrt{2H/g}$$

$$\therefore R = v_{\text{eff}} \times t = \sqrt{2gh} \sqrt{2H/g}$$

$$R^2 = 4hH \text{ or } h = \frac{R^2}{4H}$$

Hence, the correct answer is option (4)

143. (1) $x = -3t^3 + 18t^2 + 16t$

$$v = -9t^2 + 36t + 16$$

$$a = -18t + 36$$

$$a = 0 \text{ at } t = 2\text{s}$$

$$v = -9(2)^2 + 36 \times 2 + 16$$

$$v = 52 \text{ m/s}$$

144. (1) $y = 2 \sin(\omega t - kx)$

Maximum particle velocity $A\omega$

$$\text{Wave velocity} = \frac{\omega}{k}$$

$$\frac{\omega}{k} = A\omega$$

$$k = \frac{1}{A} = \frac{2\pi}{\lambda}$$

$$\lambda = 2\pi A = 4\pi \text{ cm}$$

145. (1) As refractive index, $\mu = \frac{\sin\left(\frac{A + D_{\text{max}}}{2}\right)}{\sin\frac{A}{2}}$

As angles of prism is constant, $D_{\text{min}} \propto \mu$

$\therefore \mu_b > \mu_r$, hence $D_2 > D_1$.


146. (2)

147. (3) $i = \frac{E_{\text{eq}}}{r_{\text{eq}}} = \frac{8 \times 5}{8 \times 0.2}$

$$I = 25 \text{ A}$$

$$V = E - ir$$

$$= 5 - 0.2 \times 25 = 0 \text{ V}$$



A	B	\bar{A}	\bar{B}	$\bar{A} + \bar{B}$	$\overline{\bar{A} + \bar{B}}$
0	0	1	1	1	0
0	1	1	0	1	0
1	0	0	1	1	0
1	1	0	0	0	1

149. (2) As, $U = kx^4$; $F = -\frac{dU}{dx} = -4kx^3$ (i)

In SHM, $x = r \sin \omega t$ and $\frac{d^2x}{dt^2} + \omega^2x = 0$

$$\text{Acceleration } A = \frac{d^2x}{dt^2} = -\omega^2x$$

$$\therefore F = mA = -m\omega^2x \text{(ii)}$$

From (i) and (ii);

$$4kx^3 = m\omega^2x$$

$$T = \frac{2\pi}{\omega} = \frac{2\pi}{2x} \sqrt{\frac{m}{k}} = \frac{\pi}{(r \sin \omega t)} \sqrt{\frac{m}{k}} \text{ or } T \propto \frac{1}{r}$$

150. (1) Common potential, $V = \frac{C_1V_1 + C_2V_2}{C_1 + C_2}$

$$= \frac{(4 \times 10^{-6}) \times 80 + (6 \times 10^{-6}) \times 30}{4 \times 10^{-6} + 6 \times 10^{-6}} = 50 \text{ V}$$

\therefore Energy lost by a $4 \mu\text{F}$ capacitor is


$$= \frac{1}{2}C_1V_1^2 - \frac{1}{2}C_1V^2 = \frac{1}{2}C_1(V_1^2 - V^2)$$

$$= \frac{1}{2} \times (4 \times 10^{-6}) \times \{(80)^2 - (50)^2\} = 7.8 \times 10^{-3} \text{ J} = 7.8 \text{ mJ}$$

CHEMISTRY

SECTION - A (35 Questions)

151. (2)
There is a high jump from IE_3 to IE_4 . Therefore, it is difficult to remove the 4th valence electron. So, electronic configuration of element must have three valence electrons. Hence, the answer is (2).
152. (1)
The smaller the reduction potential of a substance, the more is its reducing power.
($Y > Z > X$)
153. (3)
$$G^* = \frac{l}{a} = \frac{1.5 \text{ cm}}{0.75 \text{ cm}^2} = 2.0 \text{ cm}^{-1}$$
154. (3)
 NaBH_4 does not reduce $\text{C}=\text{C}$ bond.
155. (2)




156. (1)
 F_2 has lower bond dissociation energy than Cl_2 due to strong inter-electronic repulsions.
157. (2)
Only Xe reacts with F_2 because it has the lowest ionisation enthalpy HNO_3 .
158. (1)
 $\text{A}-\text{B}$ interaction is stronger than $\text{A}-\text{A}$ and $\text{B}-\text{B}$
159. (4)
(1)-(iii); (2)-(iv); (3)-(ii); (4)-(i)
160. (1)
The order of boiling points of the isomeric amines is as follows.
Primary amines $>$ secondary amines $>$ tertiary amines.
Boiling point \propto surface area of the molecule.
161. (2)
 $\text{H}_2\text{C}=\text{CH}_2$ Hofmann's product is the major product.
162. (4)
A-4, B-3, C-2, D-1
163. (3)
C.N. of E in complex is 6 and oxidation state is $x + 0 - 2 - 1 = 0$; $x = 3$
164. (1)
Condition for precipitation is
$$K_{\text{IP}} > K_{\text{sp}}$$


When equal volume mixed concentration become half also, for ppt,

$$[\text{Ag}^+][\text{Cl}^-] > K_{\text{sp}}$$

165. (1)
Conc. of $\text{HCl} = 0.25$ mole
Conc. of $\text{NaOH} = 0.25$ mole
Heat of neutralization of strong acid by strong base = -57.1 kJ
 $\text{HCl} + \text{NaOH} \rightarrow \text{NaCl} + \text{H}_2\text{O} - 57.1$ kJ
1 mole of HCl neutralise 1 mole of NaOH , heat evolved = 57.1 kJ
 \therefore 0.25 mole of HCl neutralise 0.25 mole of NaOH
 \therefore Heat evolved = $57.1 \times 0.25 = 14.275$ kJ
166. (4)
Since aniline gets protonated in strongly acidic medium, the lone pair of electrons are not available to produce mesomeric or electrometric effect. Thus, aniline becomes less reactive.
167. (2)



168. (4)
(1) $[\text{Co}(\text{NH}_3)_6]^{3+}$: The cobalt ion is in +3 oxidation state with $3d^6$ configuration and thus is diamagnetic octahedral complex, $[\text{Co}(\text{NH}_3)_6]^{3+}$ and has the electronic configuration represented as shown below.





- (2) $[\text{Zn}(\text{NH}_3)_6]^{2+}$: Because of $3d^{10}$ configuration no $(n-1)$ orbital is available for d^2sp^3 hybridisation and thus forms outer orbital complex. The complex is diamagnetic.
- (3) $[\text{Ti}(\text{NH}_3)_6]^{3+}$: $3d^1$ configuration and thus has one unpaired electron.
- (4) $[\text{Cr}(\text{NH}_3)_6]^{3+}$: The complex is inner orbital complex but $3d^3$ configuration has three unpaired electrons with weak as well as with strong field ligand.
169. (2)
 Zr and Hf have nearly same atomic radius as well as ionic radius. They have similar physical and chemical properties.

170. (3)
It may involve increase or decrease in temperature of the system. Systems in which such processes occur, are thermally insulated from the surroundings.

171. (4)
(1) O.N. of Cl in $\text{Cl}^- = -1$
(2) O.N. of Cl in $\text{ClO}^- = x - 2 = -1$ or $x = +1$
(3) O.N. of Cl in $\text{ClO}_2^- = x + 2 \times (-2) = -1$ or $x = +3$
(4) O.N. of Cl in $\text{ClO}_3^- = x + 3 \times (-2) = -1$ or $x = +5$

172. (3)
Iodoform test is given by compounds which have CH_3CO group.



\therefore 2-pentanone has CH_3CO group, so it gives iodoform test, while 3-pentanone does not have CH_3CO group, so it does not give iodoform test.

173. (4)
2-ethyl pent-3-en-1-ol

174. (3)
When alkaline KMnO_4 (oxidising agent) is treated with KI, iodide ion is oxidised to $2\text{IO}_3^- + 2\text{KMnO}_4 + \text{H}_2\text{O} + \text{KI} \longrightarrow 2\text{MnO}_2 + 2\text{KOH} + \text{KIO}_3$

175. (3)
 $\text{H}_2^+ = \sigma_{1s}^1$ B.O. = $\frac{1}{2}(1-0) = \frac{1}{2}$

$\text{H}_2^- = \sigma_{1s}^2 \sigma_{1s}^{*1}$ B.O. = $\frac{1}{2}(2-1) = \frac{1}{2}$

Though both have same bond order, H_2^- is less stable because it has one electron in the higher energy and antibonding molecular orbital.

176. (2)
Wavelength of yellow light = 240 nm
 $= 240 \times 10^{-9} \text{ m}$

Wave number



$$(\bar{\nu}) = \frac{1}{\lambda} = \frac{1}{240 \times 10^{-9} \text{ m}} = 4.16 \times 10^6 \text{ m}^{-1}$$

177. (3)
In case of Hydrogenic species is:
 $1s < 2s = 2p < 3s = 3p = 3d$
Hence 2s and 2p have same energy for Hydrogenic

species.

178. (1)
Aldol reaction involves an aldehyde or ketone having an α -hydrogen atom. This type of reaction occurs in presence of dilute base (i.e., dil NaOH). Only CH_3COCH_3 will give aldol reaction (Both HCHO and $\text{C}_6\text{H}_5\text{CHO}$ lack α -hydrogen).

179. (3)
Carbon and hydrogen are estimated in organic compounds by Liebig's method




Percentage of carbon and hydrogen is calculated from the weight of CO_2 and H_2 produced.

180. (1)
 $\text{NO}^\oplus > \text{NO} > \text{NO}^\ominus$


181. (1)
TTT


182. (4)
For the given reaction,



$$\text{rate} = \frac{1}{3} \frac{d[\text{Br}_2]}{dt} = -\frac{1}{5} \frac{d[\text{Br}^-]}{dt}$$


$$\therefore \frac{d[\text{Br}_2]}{dt} = -\frac{3}{5} \frac{d[\text{Br}^-]}{dt}$$

183. (2)


184. (4)

n-Butane 2-Methyl butane 2, 3-Dimethyl butane


185. (3)
Triads of group VIII (groups 8, 9 and 10) have approximately the same size e.g.
Ist triad, Fe, Co, Ni
IInd triad, Ru, Rh, Pd
IIIrd triad, Os, Ir, Pt

SECTION - B (Attempt Any 10 Questions)

186. (2)
 $\text{H}_2\text{SO}_3 < \text{H}_2\text{SO}_4 < \text{H}_2\text{S}_2\text{O}_8$

1 S = O 2 S = O 4 S = O

187. (1)

Complex of the type $[M(AA)_2X_2]$ or $[M(AA)_2XY]$ exhibit optical activity. $[CoCl_2(en)_2]^+$ forms geometrical isomers (cis and trans form). It is interesting to note that trans form does not show optical isomerism, i.e. it cannot be resolved into optical isomers. The reason is that the molecule has plane of symmetry. On the other hand, the cis-isomer is unsymmetrical and can be resolved into optical isomers.



188. (4)

$$\Lambda_m^o(NH_4Cl) + \Lambda_m^o(NaOH) - \Lambda_m^o(NaCl)$$

189. (1)

The value of K_{eq} is the measure of extent of reaction. For most stable oxide $\Rightarrow K_{eq}$ should be minimum.

190. (2)




191. (3)

Vitamin	Deficiency disease
A. Vitamin-B ₁₂	1. Pernicious anaemia
B. Vitamin-B ₆	2. Convulsions
C. Vitamin-E	3. Sterility
D. Vitamin-K	4. Haemorrhagic condition

192. (3)

Conceptual fact.



193. (4)



Here, O.N. of C decreases from +3 in $(CN)_2$ to +2 in CN^- ion and increases from +3 in $(CN)_2$ to +4 in CNO^- ion. Thus, cyanogen is simultaneously reduced to cyanide ion and oxidised to cyanate ion. It is an example of a disproportionation reaction.

194. (1)

The balanced equation involved in the titration may be given as:



$$\frac{M_1V_1}{n_1}(Cr_2O_7^{2-}) = \frac{M_2V_2}{n_2}(Fe^{2+})$$

$$\frac{0.03 \times 20}{1} = \frac{M_2 \times 15}{6}$$

$$M_2 = \frac{0.03 \times 20 \times 6}{15} = 0.24$$


$$= 24 \times 10^{-2}$$

195. (1)

All ketones in (i), (iii) and (iv) contain abstractable alpha-proton while all aldehydes do not contain alpha-hydrogen.

196. (2)

Grignard reagent forms addition product with bubbled carbon dioxide which on hydrolysis with HCl yields benzoic acid.



197. (2)

For 1 mole $Na_2CO_3 \rightarrow 1$ mole CO_2 required.

$$n_{Na_2CO_3} \Rightarrow \frac{21200}{106} = 200 \text{ moles. } [M_{Na_2CO_3} = 106 \text{ g/mole}]$$

\therefore For 200 moles $Na_2CO_3 \rightarrow 200$ moles of CO_2 required & for 1 mole $CO_2 \rightarrow 1$ mole $CaCO_3$ required.

\therefore For 200 mole $CO_2 \rightarrow 200$ mole $CaCO_3$ req.
i.e. $\Rightarrow 200 \times (100) \text{ g } CaCO_3 \text{ req. } \{m_{CaCO_3} = 100 \text{ g}\}$
 $\Rightarrow 20 \text{ Kg}$


198. (2)

$$\text{Rate} = k[A][B_2]$$

$$\text{But } K = \frac{[A]^2}{[A_2]} \text{ or } [A] = \sqrt{K \cdot [A_2]}$$

$$\text{Rate} = k \cdot \sqrt{K} \cdot \sqrt{[A_2]} [B_2] = k' [A_2]^{1/2} [B_2]$$

199. (3)



200. (1)

If both Assertion & Reason are True & the Reason is a correct explanation of the Assertion