

## ANSWER KEY & SOLUTION KEY FINAL ROUND - 06 (PCB) Dt.12.04.2024

### PHYSICS

#### SECTION - A (35 Questions)

01. (3) Since, Kiran's initial and final positions coincide.

Thus, his displacement,

$$\Delta x = x_{final} - x_{initial} = 0$$

However, corresponding path length

$$240 + 240 = 480 \text{ m}$$

Thus, the magnitude of the displacement for the given course of motion is zero but the corresponding path length is 480 m.

So, all statements are correct.

02. (4)

03. (1) When a person walks on the road, he exerts a force on floor. According to Newton's third law of motion, a reaction force exerts on the person which is being provided by the frictional force.

Thus, the frictional force helps a person to walk on a rough surface.

Thus, the statement given in option (1) is incorrect, rest are correct.

04. (3)

05. (2)

06. (3) The change in length corresponds to longitudinal strain and change in shape corresponds to shearing strain.

07. (2)

08. (3) According to parallel axes theorem,

$$I = I_{CG} + Md^2 = \frac{Ml^2}{12} + Md^2$$

$$= 300 \left[ \frac{100^2}{12} + 20^2 \right] = 3.7 \times 10^5 \text{ gm-cm}^2$$

09. (4) According to Newton's law of cooling,

$$\frac{dQ}{dt} \propto \Delta\theta$$

$$\text{But } \frac{dQ}{dt} \propto (\Delta\theta)^n \quad (\text{given})$$

$$\therefore n = 1$$

10. (4) Here,  $P_1 = 4 \text{ atm}$ ,  $T_1 = 27^\circ\text{C} = 300 \text{ K}$ ,

$$V_1 = 1500 \text{ m}^3, \\ P_2 = 2 \text{ atm}, T_2 = -3^\circ\text{C} = 270 \text{ K}, V_2 = ?$$

$$\text{As } \frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$\therefore V_2 = \frac{P_1 V_1 T_2}{T_1 P_2} = \frac{4 \times 1500 \times 270}{300 \times 2} = 2700 \text{ m}^3$$

11. (2) B.E. =  $0.042 \times 931 = 39.1 \text{ MeV}$

Number of nucleon in  ${}^7_3\text{Li}$  is 7.

$$\therefore \frac{\text{B.E.}}{\text{nucleon}} = \frac{39.1}{7} = 5.6 \text{ MeV.}$$

12. (4) As  $\sigma_1 = \sigma_2$

$$\therefore \frac{Q_1}{4\pi r_1^2} = \frac{Q_2}{4\pi r_2^2} \text{ or } \frac{Q_1}{4\pi \epsilon_0 r_1^2} = \frac{Q_2}{4\pi \epsilon_0 r_2^2}$$

$$\therefore E_1 = E_2 \text{ or } E_1/E_2 = 1 \Rightarrow E_1 : E_2 = 1 : 1$$

13. (3) Loudness of sound is given by

$$dB = 10 \log \frac{I}{I_0} \left[ \begin{array}{l} I \text{ is intensity of sound} \\ I_0 \text{ is reference intensity of sound} \end{array} \right]$$

$$\therefore 120 = 10 \log \left( \frac{I}{I_0} \right)$$

$$\Rightarrow I = 1W / m^2$$

$$\text{Also } I = \frac{P}{4\pi r^2} = \frac{2}{4\pi r^2}$$

$$\therefore r = \sqrt{\frac{2}{4\pi}} = \sqrt{\frac{1}{2\pi}} \text{ m} = 0.399 \text{ m} = 40 \text{ cm}$$

14. (2) As  $\beta = \frac{\lambda D}{d} \therefore \beta = \frac{1}{d}$ . Curve (2) is correct.

15. (3) The net flux linked with closed surfaces  $S_1, S_2, S_3$  &  $S_4$  are

$$\text{For surfaces } S_1, \phi_1 = \frac{1}{\epsilon_0} (2q)$$

$$\text{For surface } S_2, \phi_2 = \frac{1}{\epsilon_0} (q + q + q - q) = \frac{1}{\epsilon_0} 2q$$

For surface  $S_3, \phi_3 = \frac{1}{\epsilon_0}(q + q) = \frac{1}{\epsilon_0}(2q)$

For surface  $S_4, \phi_4 = \frac{1}{\epsilon_0}(8q - 2q - 4q) = \frac{1}{\epsilon_0}(2q)$

Hence,  $\phi_1 = \phi_2 = \phi_3 = \phi_4$  i.e. net electric flux is same for all surfaces.

Keep in mind, the electric field due to a charge outside ( $S_3$  and  $S_4$ ), the Gaussian surface contributes zero net flux through the surface, because as many lines due to that charge enter the surface as leave it.


16. (2)  $\frac{I_g}{I_s} = \frac{s}{R_g}$

$R_g = \frac{50 - 20}{20} \times 30\Omega \Rightarrow \frac{3}{2} \times 30\Omega = 45\Omega$

17. (2) Put  $y = 0$ ,

$0 = 12x - \frac{3}{4}x^2 \Rightarrow x = 16m$

18. (1) In the following figure, magnetic fields at O due to section 1, 2, 3 and 4 are considered as  $B_1, B_2$  and  $B_3$  and  $B_4$ , respectively.



$B_1 = B_3 = 0$

$B_2 = \frac{\mu_0}{4\pi} \cdot \frac{\pi i}{R_1} \otimes$

$B_4 = \frac{\mu_0}{4\pi} \cdot \frac{\pi i}{R_2} \odot$

As  $|B_2| > |B_4|$


So  $B_{net} = B_2 - B_4 \Rightarrow B_{net} = \frac{\mu_0 i}{4} \left( \frac{1}{R_1} - \frac{1}{R_2} \right) \otimes$

19. (1) Here,  $M = 0.4 \text{ JT}^{-1}$ ;  $B = 0.16 \text{ T}$   
For stable equilibrium, the potential energy (U) of bar magnet in the magnetic field is  
 $U = -MB = -0.4 \times 0.16 = -0.064 \text{ J}$

20. (4) Magnetic potential energy,  
 $U = 25 \text{ mJ} = 25 \times 10^{-3} \text{ J}$   
Inductance,  $L = ?$

From  $U = \frac{1}{2} LI^2$

$L = \frac{2U}{I^2} = \frac{2 \times 25 \times 10^{-3}}{(60 \times 10^{-3})^2} = \frac{500}{36} = 13.89 \text{ H}$



We have  $\omega = 120\pi$

and,  $T = \frac{2\pi}{\omega} = \frac{2\pi}{120\pi} = \frac{1}{60}$

So, req. time =  $\frac{T}{4} = \frac{1}{240} \text{ s}$

22. (3)  $X = \frac{2k^3 l^2}{m\sqrt{n}}$

The percentage error in X is given by

$\frac{\Delta X}{X} \times 100 = \left( 3 \frac{\Delta k}{k} + 2 \frac{\Delta l}{l} + \frac{\Delta m}{m} + \frac{1}{2} \frac{\Delta n}{n} \right) \times 100$

$= 3 \times 1\% + 2 \times 2\% + 3\% + \frac{1}{2} \times 4\%$


$= 3\% + 4\% + 3\% + 2\% = 12\%$

Thus, the value of X is uncertain by 12%

23. (2)  $B_0 = \frac{E_0}{c}$

E and B are in same phase.

$\hat{E} \times \hat{B} = \hat{k} \Rightarrow \hat{i} \times \hat{B} = \hat{k} \Rightarrow \hat{B} = \hat{j}$

24. (4) 

By the momentum conservation

$mv = \frac{3m}{4} v_1 \Rightarrow v_1 = \frac{4v}{3}$


25. (1) Momentum,  $p = mv = \frac{1}{2} \frac{mv \times v}{\frac{1}{2} \times v} = \frac{2KE}{v}$

If KE as well as speed are doubled, momentum p remains unchanged.

$\therefore \lambda = \frac{h}{p}$

Hence, de Broglie wavelength will remain unchanged.

26. (4)  $E = E_4 - E_3 = -\frac{13.6}{4^2} - \left( -\frac{13.6}{3^2} \right)$   
 $= -0.85 + 1.51 = 0.66 \text{ eV}$




$$a = \frac{\text{Net driving force}}{\text{Total mass in motion}} = \frac{5g}{5+4} = \frac{5g}{9}$$

28. (1) It is a system of two springs in parallel. The restoring force on the body is due to springs and not due to gravity pull.

Therefore, slope is irrelevant. Here the effective spring constant =  $k + k = 2k$ ;

Thus time period,  $T = 2\pi\sqrt{M/2k}$ .



$$n = \frac{P_2 V}{RT_2} = \frac{P_1 V}{RT_1}$$

$$\frac{P_2}{T_2} = \frac{P_1}{T_1}$$

Since  $P_2 > P_1$ , hence,  $T_2 > T_1$

30. (1) Red light and blue light have different wavelength and different frequency.

31. (3) Electron and proton have same amount of charge so they have same coulomb force. They have different acceleration because they have different masses.

32. (2)

33. (2) 
$$dq = \int_2^3 (3t^2 + 4t^3) dt = (t^3 + t^4)_2^3$$
  

$$27 + 81 - 24 = 84$$

34. (3)

35. (2) We know that,  $v_e = \sqrt{2gR}$

$$\therefore \frac{(v_e)_{P_1}}{(v_e)_{P_2}} = \frac{\sqrt{2g_1 R_1}}{\sqrt{2g_2 R_2}} = \sqrt{\frac{g_1}{g_2}} \cdot \sqrt{\frac{R_1}{R_2}} = \sqrt{kr}$$

**SECTION - B (Attempt Any 10 Questions)**

36. (4) Here,  $a = 14\text{cm}$ ,  $y = 11\text{ cm}$ ,  $V = 80\text{ m/s}$ ,  $v = ?$

$$V = \omega\sqrt{a^2 - y^2} = 2\pi v\sqrt{a^2 - y^2}$$

$$v = \frac{V}{2\pi\sqrt{a^2 - y^2}}$$

$$v = \frac{80}{2\pi\sqrt{14^2 - 11^2}} = \frac{40}{\pi \times 5\sqrt{3}} = \frac{8}{\pi \times \sqrt{3}} \text{ Hz.}$$

37. (4) The electric field on one plate due to the charge

on the other is  $E = \frac{Q}{2A\epsilon_0}$

$\therefore$  The force on one plate due to the charge on the other is

$$F = QE = Q\left(\frac{Q}{2A\epsilon_0}\right) = \frac{Q^2}{2A\epsilon_0}$$

38. (3) Impedance at resonant frequency is minimum in series LCR circuit.

$$\text{So, } Z = \sqrt{R^2 + \left(2\pi fL - \frac{1}{2\pi fC}\right)^2}$$


When frequency is increased or decreased,  $Z$  increases.

39. (1) Since, the sun is very distant,  $u$  is very large and so  $(1/u)$  is practically zero.

$$\text{So, } \frac{1}{v} + 0 = -\frac{1}{f}$$

i.e., the image of sun will be formed at the focus and will be real, inverted and diminished.

Now, as the rays from the sun subtend an angle  $\theta$  radians at the pole, hence, according to figure, we have



$$\theta = \frac{A'B'}{FP} = \frac{d}{f}$$

(where  $d$  = diameter of the image of the sun)

i.e.,  $d = \theta f$ .

40. (1)  $y = 7 \sin(7\pi t - 0.04x + \pi/3)$

Compare it with the standard equation of wave motion

$$y = r \sin\left(\frac{2\pi}{T}t - \frac{2\pi}{\lambda}x + \phi\right)$$

$$\frac{2\pi}{t} = 7\pi, T = \frac{2}{7} \text{ s, and}$$

$$\frac{2\pi}{\lambda} = 0.04, \lambda = \frac{2\pi}{0.04} = 50\pi \text{ m}$$

$$v = \frac{\lambda}{T} = \frac{50\pi}{2/7} = 175\pi \text{ m/s}$$

41. (1) SI unit of magnetic flux, induced emf coefficient of self-inductance and magnetic energy are respectively the weber, volt, henry and joule. Choice (1) is correct.

42. (2)

43. (4) Resolve the 90 N, 80 N and 70 N forces into x and y components. The line of action of 90 N, 50 N, and x-components of the 80 N and 70 N forces pass through the pivot point A, therefore they cause no rotation.

$\therefore$  The total torque about point A is

$$= (80 \sin 30^\circ) \left(\frac{L}{2}\right) - (60) \left(\frac{L}{2}\right) + (70 \cos 60^\circ)(L)$$

$$= (80) \left(\frac{1}{2}\right) \left(\frac{3}{2}\right) - (60) \left(\frac{3}{2}\right) + (70) \left(\frac{1}{2}\right) (3) = 75 \text{ N m.}$$

44. (4) When a charged particle is moving on a circular path in a magnetic field, the magnitude of velocity does not change but direction of velocity is changing every moment. Hence velocity is changing, so momentum ( $m\vec{v}$ ) is also changing.

45. (3)  $E = \frac{p^2}{2m} \Rightarrow \sqrt{E} \propto p$

$$\Rightarrow \sqrt{E} \propto \frac{1}{(1/p)}$$

46. (3)  $f = -15 \text{ cm}$  (as mirror is concave)  
 $m = -2$  (–ve sign is due to virtual image)

$$m = \frac{v}{u}$$

$$\text{or } -2 = \frac{v}{u} \Rightarrow v = -2u$$

$$\text{Also, } \frac{1}{f} = \frac{1}{v} + \frac{1}{u} = -\frac{1}{2u} + \frac{1}{u} = \frac{1}{2u}$$

$$\therefore u = \frac{f}{2} = \frac{-15}{2} = -7.5 \text{ cm.}$$

47. (1) From the principle of dimensional homogeneity  $[\alpha t] = \text{dimensionless}$

$$\therefore [\alpha] = \left[\frac{1}{t}\right] = [T^{-1}]$$

$$\text{Similarly, } [x] = \frac{[v_0]}{[\alpha]}$$

$$\therefore [v_0] = [x][\alpha] = [L][T^{-1}] = [LT^{-1}]$$

48. (1) Number significant figures 23.023 is 5.

And, that of 1.

And, that of 2.

49. (2) Gases have less viscosity.

Due to insoluble impurities like detergent surface tension decreases

50. (1) Here,  $h = 100 \text{ m}$ ,  $\Delta T = ?$

If  $m$  is mass of water, then energy converted into heat,

$$Q = mgh$$

If  $\Delta T$  is rise in temperature, then


$$Q = cm\Delta T = mgh$$

$$\Delta T = \frac{gh}{c} = \frac{10 \times 100}{4200} = 0.23^\circ \text{C.}$$

## CHEMISTRY

### SECTION - A (35 Questions)

51. (2)  
 $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O} = 2 \times 23 + 32 + 4 \times 16 + 10 \times 18$   
 $= 46 + 32 + 64 + 180 = 322 \text{ gm}$   
 $322 \text{ gm Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$  contains = 224 gm oxygen  
 $32.2 \text{ gm Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$  contains  
 $= \frac{32.2 \times 224}{322} = 22.4 \text{ gm}$
52. (4)  
 t-alkyl halide undergoes elimination.
53. (3)  
 –OR is ring activating group and ortho para directing group.
54. (1)  
 White phosphorus is soluble in  $\text{CS}_2$  whereas red phosphorus is insoluble in it.
55. (4)  
 Small hydrogen atoms can easily fit in between boron atoms but large chlorine atoms do not
56. (3)  
 The oxidation state of oxygen in  $\text{H}_2\text{O}_2$  is –1 which is an intermediate oxidation state value for oxygen hence  $\text{H}_2\text{O}_2$  can act both as reducing as well as oxidising agent.
57. (2)




$$-\frac{E_a}{R} = -5 \times 10^3 = -5000$$

$$\Rightarrow E_a = 5000 \times 8.314 = 41570 \text{ J mol}^{-1}$$

$$= 41.57 \text{ kJ mol}^{-1}$$

58. (2)




59. (1)

From Kjeldahl's method,  
Percentage of nitrogen


$$= \frac{1.4 \times N \times V}{W} = \frac{1.4 \times 0.1 \times 30}{5} = 0.84\%$$

60. (4)



Majority of four co-ordinated complexes of nickel are square planar.  
 $\text{MnX}_4^{2-}$  complex are tetrahedral.  
 $\text{F}^-$  is a weak field ligand.  $[\text{FeF}_6]^{4-}$  is thus a high spin complex with  $sp^3d^2$  hybridisation.



No unpaired electrons : diamagnetic




$$n = 5, N = \sqrt{n(n+2)} = 5.9 \text{ B.M.}$$



$$n = 4, N = \sqrt{n(n+2)} = 4.9 \text{ B.M.}$$

61 (3)



62. (4)

For a first order reaction half-life period is constant i.e. it is independent of initial concentration of the reacting species. It is related to rate constant as

$$t_{1/2} = \frac{0.693}{k}$$

63. (4)

We know that

$$\Delta H = \Delta E + P\Delta V$$

In the reactions,  $\text{H}_2 + \text{Br}_2 \rightarrow 2\text{HBr}$  there is no change in volume or  $\Delta V = 0$ .

So,  $\Delta H = \Delta E$  for this reaction.

64. (1)

(i)-(b), (ii)-(a), (iii)-(d), (iv)-(e), (v)-(c)

65. (2)

$\text{KMnO}_4$  is purple in colour due to (ligand  $\rightarrow$  metal) charge transfer phenomenon.



There is no electron present in d-orbitals of manganese in  $\text{MnO}_4^-$  (O.S. of Mn is +7).

66. (1)

Both A and R are true but R is not the correct explanation of A.

$\text{KCl}$ ,  $\text{NaCl}$  and  $\text{NH}_4\text{Cl}$  cannot be used as a salt-bridge in a cell containing silver or silver ion because they react with it to form a ppt. of  $\text{AgCl}$ .

67. (2)



$$E_{\text{cell}}^\circ = (E_{\text{R.P.}}^\circ)_{\text{cathode}} - (E_{\text{R.P.}}^\circ)_{\text{anode}}$$

$$E_{\text{cell}}^\circ = 0.34 - (-0.76) = 1.10 \text{ V}$$

$$E_{\text{cell}} = E_{\text{cell}}^\circ = 1.10 \text{ V}$$

68. (3)



(i)-(b), (ii)-(d), (iii)-(e), (iv)-(a), (v)-(c)

69. (3)

I, II and IV only

70. (3)

Melting point of Fe is more than of Mn



71. (4)

$\text{H}_2\text{O}$  is angular and  $\text{BeF}_2$  is linear

72. (3)

$$N = \frac{(10 \times 0.05) + (12 \times 0.025) + (5 \times 0.04)}{1} = 1$$

73. (2)

Among the given statements, (B) and (D) are incorrect whereas A, C and E are correct. The correct form of (B) and (D) are:

At equilibrium constant for the reverse reaction is equal to the inverse of the equilibrium constant for the forward reaction.

Equilibrium constant reflects the change in stoichiometric coefficients.


74. (2)

Acetonoxime

75. (3)

Statement-1 is false, Statement-2 is true

76. (2)



77. (2)

Methoxypropane and ethoxyethane are metamers to each other.

78. (1)

 $(\pi^*2p_x)^1$  and  $(\pi^*2p_y)^1$ 

79. (2)

$$\lambda = \frac{c}{\nu} = \frac{3 \times 10^{17}}{6 \times 10^{15}} = 50 \text{ nm}$$

80. (3)

For example for  $n = 3$  $\ell = 0, 1, 2$  (total 3 values)

81. (2)

Greater the stability lesser will be heat of hydrogenation.

Trans-2-butene more stable, less heat of hydrogenation.

82. (3)

Electrophilic addition reaction involves rearrangement of carbocation.

83. (3)

Conceptual fact.

84. (2)

The second ionisation potential of Mg is greater than the second ionisation potential of Na.

85. (4)

Multiple proportions

### SECTION - B (Attempt Any 10 Questions)


86. (3)  $2\text{BrO}_3^- + 12\text{H}^+ + 10\text{Br}^- \rightarrow 6\text{Br}_2 + 6\text{H}_2\text{O}$ 

10 mole  $e^-$  required for formation of 6 moles of  $\text{Br}_2$


$$\therefore \text{n-factor of } \text{Br}_2 = \frac{10}{6} = \frac{5}{3}$$

$$\text{eq.wt.} = \frac{\text{mol.wt.}}{n} = \frac{m}{5/3} = \frac{3M}{5}$$

87. (2)



$$\Delta H_1 = -55.84 \times 2 = -111.68 \text{ kJ/mol}$$




$$\Delta H_2 = -106.68 \text{ kJ/mol}$$

$$\Delta H_{\text{ionisation}} = 111.68 - 106.68 = 5 \text{ kJ/mol}$$

88. (4)

Cannizzaro's reaction

89. (1)



90. (4)

2 and 4

91. (2)

A-II, B-III, C-IV, D-I

92. (1)

$$\pi_1 = \pi_2$$

$$C_1 = C_2$$


$$\frac{5.12}{342} = \frac{0.9}{M}$$

$$M = 60$$

93. (4)

Due to resonance,  $\text{C}_2 - \text{C}_3$  bond is little shorter than  $\text{C} - \text{C}$  single bond length of  $1.54 \text{ \AA}$  in ethane. So the most appropriate value is  $1.46 \text{ \AA}$ .

94. (2)

Let P is initial pressure of  $\text{NO}_2$ At eqm  $P - 2x$        $2x$        $x$ as per given  $x = 0.25$ 

$$K_p = \frac{(2x)^2 (x)}{(P - 2x)^2}$$


$$\Rightarrow 156.25 = \frac{(0.5)^2 (0.25)}{P_{\text{NO}_2}^2}$$

$$\Rightarrow P_{\text{NO}_2} = 0.02$$

95. (3)

(P)-(2); (Q)-(1); (R)-(4); (S)-(3)

96. (2)



$$\text{B.O.} = \frac{5}{4} = 1.25$$

Due to resonance  $-3$  charge is distributed in all the four oxygen atoms hence each oxygen has

$$\frac{-3}{4} = -0.75 \text{ charge.}$$

97. (3)  
If (A) is right but (R) is wrong.  
In allenes terminal double bonded carbons must be connected to two different groups.
98. (2)  
A-(t), B-(p), C-(s), D-(q), E-(r)
99. (1)  
Due to electronegativity difference, the stability of interhalogen compounds follows following order:  
 $\text{IF}_3 > \text{BrF}_3 > \text{ClF}_3$
100. (2)  
Moles of  $\text{CoCl}_3 \cdot 6\text{NH}_3$   
$$= \frac{2.675}{267.5} = \frac{1}{100} = 0.01 \text{ mole}$$
  
Moles of  $\text{AgCl} = \frac{4.78}{143.5} = \frac{3}{100} \text{ mole} = 0.03 \text{ mole}$   
0.01 mole of compound gives moles of  $\text{AgCl} = 0.03 \text{ mole}$  So, 1 mole of compound gives  
moles of  $\text{AgCl} = \frac{0.03}{0.01} = 3 \text{ mole}$ . So structural formula of compound having  $3\text{Cl}^-$  ions out side of coordination sphere, so formula is  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$ .

## BOTANY

### Section - A (35 Questions)

101. (2) (NCERT XII, Pg 91, Fig 5.15)  
102. (2) (NCERT XII, Pg 111, Para 1, Line 2)  
103. (4) (NCERT XII, Pg 109, Based on Transcription process)  
104. (3) (NCERT XII, Pg 129, Para 1, Line 2)  
105. (2) (11th Para 10.4.1, Page no.168)  
106. (1) (NCERT XII, Pg 89, Mendelian disorder)  
107. (3) (NCERT XII, Pg 92, Down's Syndrome-3<sup>rd</sup> line)  
108. (3) (11<sup>th</sup> NCERT PK, page no.28, fig.3.2(d))  
109. (2) (11<sup>th</sup> NCERT PK, 1MMC=4spores means  $10\text{MMC} \times 4 = 40$  Spores, 1spore = 1Protonema = 10 leafy stage.)

110. (3) (12<sup>TH</sup> NCERT page no.248 Fig.14.4(a),14.4(c))  
111. (3) (11th Para 8.5.10, Figure 8.13, Page no.139)  
112. (4) [NCERT XI, New added family]  
113. (2) (NCERT XI Pg.227, 2<sup>nd</sup> Para)  
114. (3) (NCERT XI Pg.235, 14.4, 2<sup>hd</sup> Para, 2<sup>nd</sup> line)  
115. (4) (NCERT XII, Pg 80, based on Law of Independent Assortment)  
116. (2) (NCERT 12<sup>th</sup>, Page no- 31, 2<sup>nd</sup> paragraph, Line no- 15-17)  
117. (2) (NCERT 12<sup>th</sup>, Page no- 36, Paragraph- 2.4.3, Line no- 12,13)  
118. (1) (NCERT 11<sup>th</sup>, Page no- 26, 2<sup>nd</sup> paragraph, Line no- 21, 22)  
119. (4) (NCERT 11<sup>th</sup>, Page no- 7, 1<sup>st</sup> paragraph, Line no- 1-4)  
120. (4) (NCERT XI page no. 214, fig. 13.7)  
121. (2) (NCERT XI page no. 215, sub-topic 13.7, 3<sup>rd</sup> paragraph, last 2 lines)  
122. (4) (NCERT XII, Pg 117, Fig- 5.14)  
123. (2) (NCERT XII, Pg 115, Para 3, Line 5)  
124. (1) (NCERT XII, Pg 97, Based on Chargaff's rule)  
125. (3) (11th Para 8.5.10, Page no.139)  
126. (3) (11th Para 8.4.2, Page no.129)  
127. (1) (11th Para 10.2, Page no.164,165,166)  
128. (1) [NCERT class XI, Page no. 249 (First paragraph) and 250 (Point 15.4.3.4 (Third paragraph))]  
129. (1) [NCERT XI, Page No.75; Sub-topic 5.5.1.3 & 5.5.1.4]  
130. (3) [NCERT XI, Page No. 80; Sub-topic 5.9.2]  
131. (4) [NCERT class XI, Page no. 87, Point 6.1.2.2 (Last line), 88 (Line no. 01)  
132. (3) (NCERT Page no- 23, 2<sup>nd</sup> Paragraph, Concept based)  
133. (4) (NCERT 11<sup>th</sup>, Page no- 22, Paragraph- 2.3, Line no- 18- 22)  
134. (4) [NCERT class XI, Page no. 91, First paragraph- Last line]  
135. (2) (11<sup>th</sup> NCERT Page no.32, Concept)

### SECTION - B (Attempt Any 10 Questions)

136. (3) (NCERT Page no- 27, 1<sup>st</sup> Paragraph, Line no- 20,21)  
137. (3) (NCERT XI page no. 213, fig. 13.6 and 2<sup>nd</sup> paragraph)  
138. (3) [NCERT XI, Page No. 67, 71 & 77; Sub-topic 5.1.1; 5.3.3; 5.4; 5.7.2]  
139. (4) (NCERT 11<sup>th</sup>, Page no- 7, Last paragraph, Line no- 33-37)

140. (2) (NCERT 11<sup>th</sup>, Page no- 23, Paragraph- 2.3.1, Line no- 3,4)
141. (3) (12<sup>th</sup> NCERT Page no.244 ,14.3,3<sup>rd</sup> para)
142. (3) (NCERT XII ,Pg 91, Phenylketonuria)
143. (1) (11<sup>th</sup> Para 8.4.1, Page no.129)
144. (3) (NCERT XII, Pg 112, Para 1, Line 4)
145. (3) (11<sup>th</sup> Para 10.4, concept based- Page no.167)
146. (1) [NCERT class XI, Page no. 94, Line number- 06-09]
147. (1) [NCERT class XI,, Page 250, Point 15.4.3.4 (First paragraph)]
148. (1) (NCERT XI Pg.229, Last Para, 1<sup>st</sup> line)
149. (3) (11<sup>th</sup> NCERT Conceptual)
150. (4) (NCERT Page no 20, Paragraph- 2.2, line no- 19,20)

## ZOOLOGY

### Section - A (35 Questions)

151. (1) (NCERT XI Page No. 52; phylum annelida)
152. (1) (NCERT XI Page No. 49, Phylum-porifera)
153. (3) (12<sup>th</sup> NCERT Page no.263 1<sup>st</sup> para)
154. (4) (12<sup>th</sup> Para 10.1, Page no.181)
155. (4) (12<sup>th</sup> Para 10.1, Page no.181)
156. (3) (NCERT XI NCERT conceptual)
157. (3) ( 12<sup>th</sup> NCERT page no.219,13.1.2 (i), 225 2<sup>nd</sup> paras)
158. (3) (NCERT11<sup>th</sup>, page no 102, para 3)
159. (3) (NCERT11<sup>th</sup>, page no 116, para 2, line 6)
160. (1) (NCERT 12<sup>th</sup> page no 52,53)
161. (4) (NCERT Pg. 269)
162. (3) ( NCERT Pg. No. 286)
163. (4) (12<sup>th</sup> NCERT Page no.231, conceptual)
164. (2) (NCERT 12<sup>th</sup>, page no 44, last para)
165. (3) (NCERT 12<sup>th</sup>, page no 46, para 2)
166. (1) (NCERT12<sup>th</sup> page no 46, para 2)
167. (3) (NCERT 12<sup>th</sup> page no 64, para 3)
168. (4) (NCERT Pg. No. 157)
169. (4) (NCERT Pg. No.157)
170. (3) [NCERT P.No.308 Last Para, 304 Last Para, P.No. 305 10<sup>th</sup> Line]

171. (1) [NCERT P.No.310 2<sup>nd</sup> para, 15<sup>th</sup> line ]
172. (4) [NCERT P. No.321 1<sup>st</sup> Line]
173. (3) (NCERT XI<sup>th</sup> Page No. 270)
174. (4) (NCERT Pg. No. 149, 150)
175. (2) (NCERT Pg. No. 283)
176. (3) (NCERT 11<sup>th</sup>, Page no- 146, Paragraph- 9.3, Concept based)
177. (3) [NCERT P.No.210,12.2 1<sup>st</sup> para]
178. (3) [NCERT P.No.317, Generation and conduction of Nerve Impulse ]
179. (2) (NCERT 12<sup>th</sup>, Page no- 142, Summary)
180. (3) (NCERT 12<sup>th</sup>, Page no- 134, Figure- 7.7 concept based)
181. (2) (NCERT Page no- 137, Paragraph-3, First 2 lines)
182. (1) [NCERT P.No.199 Fig 11.4,]
183. (1) [NCERT P.No.208 GMO Points Applied,]
184. (3) (NCERT 11<sup>th</sup>, Page no- 144, 2<sup>nd</sup> paragraph, ,Line no- 9, 10)
185. (1) (NCERT XI Page No. 293; 4<sup>th</sup> line of 5<sup>th</sup> paragraph.)

### SECTION - B (Attempt Any 10 Questions)

186. (4) [NCERT P.No.208, GMO Points Applied]
187. (4) (NCERT Pg. No. 148)
188. (2) (NCERT Pg. No. 157)
189. (2) (NCERT XI Page No. 333, Last 2 lines of 1<sup>st</sup> paragraph)
190. (1) (NCERT XI Page No. 48, 4.1.6)
191. (1) [NCERT P.No.312, Synovial Joints 4<sup>th</sup> Line]
192. (3) (12<sup>th</sup> NCERT Page no.233, (i), concept)
193. (3) (NCERT 12<sup>th</sup> page no 59, para 2)
194. (1) (NCERT XI Page No. 333, 6<sup>th</sup> line of 2<sup>nd</sup> paragraph)
195. (4) (NCERT 11<sup>th</sup> page no 114, para 3)
196. (3) (NCERT 12<sup>th</sup>, Page no- 135, 3<sup>rd</sup> paragraph, Line no- 1-10)
197. (2) (NCERT 11<sup>th</sup>, Page no-147, Table-9.4)
198. (3) [NCERT P.No.319, First Para ]
199. (4) [NCERT P.No.310 12<sup>th</sup> Line ]
200. (3) (12<sup>th</sup> Para10.3 based /Page no. 184)