


## ANSWER KEY & SOLUTION KEY FINAL ROUND - 11 (PCB) Dt.20.04.2024

### PHYSICS

#### SECTION - A (35 Questions)

01. (2) Smaller gulab jamuns are having least surface area hence, they will be heated first. As in case of smaller gulab jamun heat radiates will be less. Similarly, smaller pizzas are heated before bigger ones because they are of small surface areas.
02. (3) As the body is falling freely under gravity, the potential energy decreases and kinetic energy increases but total mechanical energy (PE + KE) of the body and earth system will be constant as external force on the system is zero.
03. (2)
04. (4) Given,  $A = 1.0\text{m} \pm 0.2\text{m}$ ,  $B = 2.0 \pm 0.2\text{m}$ .  
 Let,  $Y = \sqrt{AB} = \sqrt{(1.0)(2.0)} = 1.414\text{m}$   
 Rounding off to two significant digit  $Y = 1.4\text{m}$   
 $\frac{\Delta Y}{Y} = \frac{1}{2} \left[ \frac{\Delta A}{A} + \frac{\Delta B}{B} \right] = \left[ \frac{0.2}{1.0} + \frac{0.2}{2.0} \right] = \frac{0.6}{2 \times 2.0}$   
 $\Rightarrow \Delta Y = \frac{0.6}{2 \times 2.0} = \frac{0.6 \times 1.4}{2 \times 2.0} = 0.212$   
 Rounding off to one significant digit  $\Delta Y = 0.2\text{m}$   
 Thus, correct value for  
 $= \sqrt{AD} = r + \Delta r = 1.4 \pm 0.2\text{m}$ .
05. (1)
06. (2) In the given diagram, when the small piece Q removed and glued to the centre of the plate, the mass comes closer to the z-axis, hence, moment of inertia decreases.
07. (4)  $\lambda = \frac{h}{p} = \frac{h}{m\sqrt{gH}}$ .
08. (1) As the lift is coming in downward direction displacement will be negative. We have to see whether the motion is acceleration or retarding.  
 We know that due to downward motion displacement will be negative. When the lift reaches 4th floor is about to stop hence, motion is retarding in nature hence,  $x < 0$ ,  $a > 0$



As displacement is in negative direction, velocity will also be negative i.e.,  $v < 0$ .

This can be shown on the adjacent graph.

09. (2)
10. (1) Let equation of an SHM is represented by  
 $y = a \sin \omega t$   
 $v = \frac{dy}{dt} = a\omega \cos \omega t$   
 $\Rightarrow (v)_{\max} = a\omega = 30 \dots\dots\dots(i)$   
 Acceleration  $(A) = \frac{d^2x}{dt^2} = a\omega^2 \sin \omega t$   
 $A_{\max} = a\omega^2 = 60 \dots\dots\dots(ii)$   
 Eqs. (i) and (ii), we get  $\omega(a\omega) = 60 \Rightarrow \omega(30) = 60$   
 $\Rightarrow \omega = 2\text{rad/sec}$   
 $\Rightarrow \frac{2\pi}{T} = 2\text{rad/sec} \Rightarrow T = \pi \text{ sec.}$
11. (2)
12. (2) Given, mass =  $m = 5 \text{ kg}$   
 acting force =  $\vec{F} = (-3\hat{i} + 4\hat{j})\text{N}$ .  
 Initial velocity at  $t = 0$ ,  $v = (6\hat{i} - 12\hat{j}) \text{ m/s}$   
 Retardation,  $\hat{a} = \frac{F}{m} = \left( -\frac{3\hat{i}}{5} + \frac{4\hat{j}}{5} \right) \text{ m/s}^2$   
 As final velocity is along Y-axis only, its x-component must be zero.  
 Form  $v = u + at$ , for X-component only,  
 $0 = 6\hat{i} - \frac{3\hat{i}}{5}t \Rightarrow t = \frac{5 \times 6}{3} = 10\text{s}.$
13. (3)



$$= \left( \frac{3.75/2}{2.5/2} \right)^2 = \left( \frac{3.75}{2.5} \right)^2 = \frac{9}{4} \left[ \begin{matrix} r_2 = \frac{d_2}{2} \\ r_1 = \frac{d_1}{2} \end{matrix} \right]$$

34. (4) The angle between velocity and magnetic field will be zero. Hence electron will not experience any force.

35. (2)  $\beta_{central} = \frac{2\lambda D}{a} \Rightarrow a = \frac{2\lambda D}{\beta_{central}}$

**SECTION - B (Attempt Any 10 Questions)**

36. (4) We know for an ideal gas,  $pV = nRT$  where,  $n$  = number of moles,  $p$  = Pressure,  $V$  = Volume  $R$  = Gas constant,  $T$  = temperature

$$= \frac{pV}{RT}$$

As number of moles of the gas remains fixed, hence, we can write

$$\frac{p_1 V_1}{RT_1} = \frac{p_2 V_2}{RT_2} \Rightarrow p_2 = (p_1 V_1) \frac{T_2}{V_2 T_1}$$

$$= \frac{(p)(V)(1.1T)}{(1.05)V(T)} [p_1 = p, V_2 = 1.05 V \text{ and } T_2 = 1.1T]$$


$$= p \times \frac{1.1}{1.05}$$

$$= p \times (1.0476) = 1.05 p.$$

Hence, final pressure  $p_2$  lies between  $p$  and  $1.1 p$ .

37. (1)

38. (3) Let  $A$  moves up the plane frictional force on  $A$  will be downward as shown.



When  $A$  just starts moving up

$$mg \sin \theta_1 + f = mg \sin \theta_2$$

$$\Rightarrow mg \sin \theta_1 + \mu mg \cos \theta_1 = mg \sin \theta_2$$

$$\Rightarrow \mu = \frac{\sin \theta_2 - \sin \theta_1}{\cos \theta_1}$$

When  $A$  moves upwards

$$f = mg \sin \theta_2 - mg \sin \theta_1 > 0$$

$$\Rightarrow \sin \theta_2 > \sin \theta_1 \Rightarrow \theta_2 > \theta_1.$$

39. (3)

40. (4) Given, fundamental quantities are momentum ( $p$ ), area ( $A$ ) and time ( $T$ ).

We can write energy  $E$  as

$$E \propto p^a A^b T^c$$

$$E = kp^a A^b T^c$$

where  $k$  is dimensionless constant of proportionality.

Dimensions of

$$E = [E] = [ML^2T^{-2}] \text{ and } [p] = [MLT^{-1}]$$

$$[A] = [L^2]$$

$$[T] = [T]$$

$$[E] = [k][p]^a [A]^b [T]^c$$

$$= M^a L^{2b+a} T^{-a+c}$$

By principle of homogeneity of dimensions,

$$a = 1, 2b + a = 2$$


$$\Rightarrow 2b + 1 = 2$$

$$\Rightarrow b = 1/2 - a + c = -2$$

$$c = -2 + a = -2 + 1 = -1$$

$$\text{Hence, } E = pA^{1/2}T^{-1}.$$

41. (4) Let the radius of the sphere is  $R$ . As the temperature increases radius of the sphere increases as shown.



$$\text{Original volume } V_0 = \frac{4}{3} \pi R^3$$

Coefficient of linear expansion =  $\alpha$

$$\therefore \text{Coefficient of volume expansion} = 3\alpha$$

$$\therefore \frac{1dV}{VdT} = 3\alpha \Rightarrow dV = 3V\alpha dt \Rightarrow 4\pi R^3 \alpha \Delta T$$

= increase in the volume.


42. (3) Consider the adjacent diagram when  $M_1$  comes in contact with the spring,  $M_1$  is retarded by the spring force and  $M_2$  is accelerated by the spring force.

(1) The spring will continue to compress until the two blocks acquire common velocity.

(2) As surface are frictionless momentum of the system will be conserved

(3) If spring is massless whole energy of  $M_1$  will be imparted to  $M_2$  and  $M_1$  will be at rest, then

(4) Collision is inelastic, even if friction is not involved.



43. (1) Given, mass =  $m = 5 \text{ kg}$ , Radius =  $1 \text{ m} = R$   
 Revolution per minute  $\omega = 300 \text{ rev/min}$   
 $= (300 \times 2\pi) \text{ rad/min}$   
 $= (300 \times 2 \times 3.14) \text{ rad/min}$   
 $= (300 \times 2 \times 3.14) \text{ rad/60s}$   
 $= \frac{300 \times 2 \times 3.14}{60} \text{ rad/s} = 10\pi \text{ rad/s}$

$\Rightarrow$  Linear speed =  $v = \omega R$   
 $= \left( \frac{300 \times 2\pi}{60} \right) (1) = 10\pi \text{ m/s}$

$KE = \frac{1}{2} m v^2$

$KE = \frac{1}{2} \times 5 \times (10\pi)^2$

$KE = 100\pi^2 \times 5 \times \frac{1}{2} = 250\pi^2 J.$

44. (2) We know that Young's modulus

$Y = \frac{\text{Stress}}{\text{Strain}} = \frac{F/A}{\Delta L/L} = \frac{F}{A} \times \frac{L}{\Delta L}$

$= \frac{F}{\pi(D/2)^2} \times \frac{L}{\Delta L} = \frac{4FL}{\pi D^2 \Delta L}$

$\Rightarrow D^2 = \frac{4FL}{\pi \Delta L Y} \Rightarrow D = \sqrt{\frac{4FL}{\pi \Delta L Y}}$

As  $F$  and  $\frac{L}{\Delta L}$  are constants.

Hence,  $D \propto \sqrt{\frac{1}{Y}}$

Now, we can find ratio as  $\frac{D_{\text{copper}}}{D_{\text{iron}}} = \sqrt{\frac{Y_{\text{iron}}}{Y_{\text{copper}}}}$ .

45. (1) As no external torque acts on the system, angular momentum should be conserved.

Hence  $I\omega = \text{constant} \dots\dots\dots(i)$

where,  $I$  is moment of inertia of the system and  $\omega$  is angular velocity of the system.

$I_1\omega_1 = I_2\omega_2$

(where  $\omega_1$  and  $\omega_2$  are angular velocities before and after jumping)

$\Rightarrow I\omega = \frac{I}{2} \times \omega_2$

(as mass reduced to half, hence, moment of inertia also reduced to half)

$\Rightarrow \omega_2 = 2\omega.$

46. (4) Gravitational mass of proton is equivalent to its inertial mass and is independent of presence neighbouring heavy objects.

47. (2)

48. (2) Mass  $m = 2.5 \text{ kg}$ ,  $\mu = \text{mass per unit length}$

$= \frac{m}{l} = \frac{2.5 \text{ kg}}{20} = \frac{125}{10} = 0.125 \text{ kg/m}$

Speed  $v = \sqrt{\frac{T}{\mu}} = \sqrt{\frac{200}{0.125}}$  [speed of transverse waves in any string]


$l = v \times t \Rightarrow 20 = \sqrt{\frac{200}{0.125}} \times t$

$t = 20 \times \sqrt{\frac{25 \times 5}{2 \times 10^5}}$

$= 20 \times \sqrt{25 \times \frac{1}{0.4 \times 10^5}} = 20 \times 5 \sqrt{\frac{1}{4 \times 10^4}}$

$= \frac{20 \times 5}{2 \times 10^2} = \frac{1}{2} = 0.5.$

49. (3) Consider the diagram in which a liquid column oscillates. In this case, restoring force acts on the liquid due to gravity. Acceleration of the liquid column, can be calculated in terms of restoring force.



Restoring force  $f = \text{Weight of liquid column of height } 2y$

$\Rightarrow f = -(A \times 2y \times \rho) \times g = -2A\rho gy$

[ $\because \eta = \rho v$ ]

$\Rightarrow f \propto -y \Rightarrow$  Motion is SHM with force constant

$k = 2A\rho g$

$\Rightarrow$  Time period

$T = 2\pi \sqrt{\frac{m}{k}} = 2\pi \sqrt{\frac{A \times 2h \times \rho}{2A\rho g}} = 2\pi \sqrt{\frac{h}{g}}$

$$T = 2\pi\sqrt{\frac{l}{g}}, \text{ where } l = h$$

Which is independent of the density of the liquid.

50. (3)

## CHEMISTRY

### SECTION - A (35 Questions)

51. (3)  $O_2, B_2$

52. (1) (i) 4p, (ii) 4s, (iii) 3d, (iv) 3p


Their order of increasing energy

(iv) < (ii) < (iii) < (i)


53. (2) For exothermic reactions,  $H_R > H_P$ , hence  $\Delta H$  is negative

54. (3) Iodoform test is used to test presence of  $-COCH_3$  group which is converted into  $-COOH$  group.

The reaction is shown as



55. (4) Source of nitrogen in Gabriel phthalimide synthesis is potassium phthalimide.



56. (4) Ionisation enthalpy of an atom is always negative.

57. (1) In Wilkinson's catalyst (a homogenous catalyst),  $(\text{Ph}_3\text{P})_3\text{RhCl}$ , Rh is  $dsp^2$  hybridised giving a square planar shape to the compound and is in +1 oxidation state.


58. (4) (A)  $\rightarrow$  (iii), (B)  $\rightarrow$  (iv), (C)  $\rightarrow$  (i), (D)  $\rightarrow$  (ii)

59. (1) I and IV statements are correct.

60. (4) In option (4) a carbon with double bond has two same functional groups ( $\text{CH}_3$ ) attached. The rotation around carbon will not produce a new compound. Hence, geometrical isomerism is not possible.

61. (2) +I-effect decreases the stability of carbon anion. Since,  $(\text{CH}_3)$  group has +I-effect, therefore, it intensifies the negative charge and hence destabilises (A) relative to (B).  $sp$  hybridised

carbanion is more stabilised than  $sp^3$



Hence,  $B > A > C$


62. (1) +800 & +640

63. (4) The transition metals exhibit higher enthalpies of atomisation due to strong interatomic interaction arises because of having large number of unpaired electrons in their atoms.

64. (1)

Species	Conjugate acid	Conjugate base
$\text{HCO}_3^-$	$\text{H}_2\text{CO}_3$	$\text{CO}_3^{2-}$
$\text{HSO}_4^-$	$\text{H}_2\text{SO}_4$	$\text{SO}_4^{2-}$
$\text{NH}_3$	$\text{NH}_4^+$	$\text{NH}_2^-$
$\text{H}_2\text{O}$	$\text{H}_3\text{O}^+$	$\text{OH}^-$

65. (3) Sucrose (cane sugar) is a disaccharide. One molecule of sucrose on hydrolysis gives one molecule of glucose and one molecule of fructose.



Note:- Sucrose is a dextro-rotatory sugar on hydrolysis produces a laevorotatory mixture. So, known as invert sugar. Sucrose is a non-reducing sugar while maltose and lactose are reducing sugar.

66. (4)  $\text{Br}_2 \rightarrow \text{BrO}_3^-$

For  $\text{Br}_2$ , oxidation number = 0

For  $\text{BrO}_3^-$ , oxidation number,  $x + (-6) = -1$   
 $\Rightarrow x = +5$

67. (1) Electronegativity of Cl, Br, C and Mg follows the order  $\text{Cl} > \text{Br} > \text{C} > \text{Mg}$

\*  $\text{CH}_3 \rightarrow \text{CH}_2 \rightarrow \text{Cl}$  (-I-effect)

\*  $\text{CH}_3 \leftarrow \text{CH}_2 \leftarrow \text{Mg}^+\text{Cl}^-$

\*  $\text{CH}_3 \rightarrow \text{CH}_2 \rightarrow \text{Br}$  (-I-effect)

\*  $\text{CH}_3-\text{CH}_2-\text{CH}_3$  (+I-effect)

-I-effect of  $\text{Cl} > \text{Br}$ .

Hence,  $\text{CH}_3-\text{CH}_2-\text{Cl}$  has the greatest positive charge.

68. (2)  $\overset{+5}{\text{HClO}_3} \rightarrow \overset{+7}{\text{HClO}_4}$  (Oxidation)

$\overset{-3}{\text{NH}_4^+} \rightarrow \overset{-3}{\text{NH}_3}$  (None)



$\overset{-4}{\text{NO}_2} \rightarrow \overset{+4}{\text{N}_2\text{O}_2}$  (None)

$\overset{+4}{\text{HSO}_3^-} \rightarrow \overset{+6}{\text{SO}_4^{2-}}$  (Oxidation)

$\overset{-1}{\text{H}_2\text{O}_2} \rightarrow \overset{-2}{\text{H}_2\text{O}}$  (Reduction)

69. (1) (p) At anode: oxidation takes place.


- (q) At cathode : reduction takes place.  
 (r) Salt bridge for migration of ions  
 (s) Electrons flow from anode to cathode.  
 (t) Current flows from cathode to anode.
70. (2) A and D  
 71. (3) Hexane + CO<sub>2</sub>  
 72. (4) Hypo solution (Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>) is used in photography to remove the unaffected AgBr in the form of soluble complex.  

$$\text{AgBr} + \text{Na}_2\text{S}_2\text{O}_3 \rightarrow \text{Na}_3[\text{Ag}(\text{S}_2\text{O}_3)_2] + 2\text{NaBr}$$
 Sodium argentothiosulphate
73. (3) Solubility changes with temperature. A plant cell shrinks in hypertonic solution. Relative lowering of vapour pressure is a colligative property.
74. (1)
- | Hydrocarbons   | Boiling point  |   |
|----------------|----------------|---|
| A. n-pentane   | 309 K due to   | no branch   |
| B. iso-pentane | 301 K due to   | one branch  |
|                |                | $\text{CH}_3-\text{CH}_2-\overset{\text{CH}_3}{\text{CH}}-\text{CH}_3$            |
| C. neo-pentane | 282.5 K due to | two branches  |
|                |                | $\text{CH}_3-\overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}}-\text{CH}_3$ |
75. (3) KCl undergoes dissociation in solution, hence observed molar mass will be lower. Experimentally determined molar mass can be higher or lower depending upon whether solute undergoes dissociation or association.
76. (4) Lanthanum hydroxide is more basic than lutetium hydroxide since the basicity decreases in the period. Atomic radius of Zr and Hf are same because of lanthanide contraction
77. (2) Cations, total charge, anions, ions, oxidation
78. (1) Homolysis process require sunlight, U.V light or high temp. to form Cl radicals.
79. (1) X = , Y = 
80. (2) A-(iii), B-(iv), C-(ii), D-(i)
81. (1) Molarity  

$$= \frac{\text{Wt. of solute}}{\text{Mol. wt. of solute}} \times \frac{1000}{\text{Volume of soln. (mL)}}$$

$$= \frac{49}{98} \times \frac{1000}{250} = 2\text{M}$$
82. (2) In <sup>88</sup><sub>38</sub>Sr, Atomic number = No. of protons  
 = No. of electrons = 38.  
 Atomic mass = 88  
 Number of neutrons = 88 - 38 = 50

83. (3) Ethanal (CH<sub>3</sub>CHO) is an oxidised product of ethanol. Pyridinium chlorochromate (C<sub>5</sub>H<sub>5</sub><sup>+</sup>NHClCrO<sub>3</sub><sup>-</sup>) oxidises primary alcohols to aldehydes. Strong oxidising agents such as KMnO<sub>4</sub> are used for getting carboxylic acid from alcohols. The oxidation process can be stopped at the reagent of pyridinium chlorochromate and pyridinium dichromate [(C<sub>5</sub>H<sub>5</sub>NH<sub>2</sub>)<sup>2+</sup>Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup>] in anhydrous medium are used as the oxidising agent. So, the correct option is (3).




84. (3) Assertion is correct but reason is wrong statement.

Correct Reason This reaction proceeds through S<sub>N</sub>2 mechanism, in which -OH ion attacks at 180° to the halogen atom of 2-bromooctane which leads to the inversion of configuration.


85. (1) Triangular planar

### SECTION - B (Attempt Any 10 Questions)

86. (2) Ph-CH<sub>2</sub>-MgI  $\xrightarrow{\text{H}_2\text{O}}$  Ph-CH<sub>3</sub> + Mg(OH)I
87. (3) Iron coated with zinc does not get rusted even if cracks appear on the surface because Zn will take part in redox reaction not Fe as Zn is more reactive than Fe. If iron is coated with tin and cracks appear on the surface, Fe will take part in redox reaction because Sn is less reactive than Fe.
88. (1) Both A and R are correct and R is the correct explanation of A.



Pent-1-ene



Pent-2-ene

When two or more compounds differ in the position of substituent atom or functional group on the carbon skeleton then it is position isomerism. Double bond is a functional group whose position varies.

89. (2)  $K = k_1 \times k_2 = (6.8 \times 10^{-3}) \times (1.6 \times 10^{-3})$   
 $= 1.08 \times 10^{-5}$
90. (1)  $\text{MgCl}_2 \rightleftharpoons \text{Mg}^{2+} + 2\text{Cl}^-$ ,  $i = 3$   
 $\Delta T_b = iK_b m = 3 \times 0.52 \times \frac{9.43}{94.3 \times 1} = 0.156$
91. (4) The correct increasing order of basic strength is as follows  
 II < IV < I < III

Greater the electron density towards ring, greater will be its basic strength. Electron withdrawing group decreases basic strength while electron donating group increases basic strength.


92. (1) A and R both are correct, and R is the correct explanation of A.

93. (4)

Element	%	No. of moles	Molar ratio	Whole no. ratio
P	27.3	$27.3/12 = 2.27$	1	1
Q	72.7	$72.7/16 = 4.54$	2	2

Empirical formula =  $PQ_2$

94. (2) Chemical reaction can be shown as



Thus,  $\text{CH}_3\text{CH}-\text{OH}$  and  $\text{CH}_3-\text{CH}_2-\text{CH}_2\text{OH}$  are

positional isomers.


Hence, option (2) is correct.

95. (1)  $sp^3$ , tetrahedral

96. (2)  $k = \frac{2.303}{(t_2 - t_1)} \log \frac{[A_1]}{[A_2]}$

$$k = \frac{2.303}{(1600 - 800)} \log \frac{1.45}{0.88} = \frac{2.303}{800} \times 0.2169$$

$$= 6.24 \times 10^{-4} \text{ s}^{-1}$$



98. (2) (ii), (iii) & (iv)

99. (1)  $\text{NH}_2-\text{Hg}-\text{O}-\text{Hg}-\text{I}$

100. (3)  $\frac{0.693}{t_{1/2}} = \frac{2.303}{t} \log \frac{a}{a-x}$

$$\frac{0.693}{10} = \frac{2.303}{100} \log \frac{100}{100-x}$$

$$\Rightarrow x = 99.9\%$$

## BOTANY

### Section - A (35 Questions)

101. (3) (Exemplar 12<sup>th</sup> Question No.07)

102. (2) (NCERT 11<sup>th</sup>, Biological Classification, Exemplar)

103. (3) (NCERT 11<sup>th</sup>, Biological Classification, Summary)

104. (1) (NCERT 11<sup>th</sup>, Biological Classification, Exercise based)

105. (4) (11<sup>th</sup> NCERT EXEMPLAR/ PAGE NO. 50)

106. (1) [NCERT Exemplar problems - XI page No. 22; Q.No.6]

107. (1) (NCERT XII, Pg 117, Para 2, Line 1)

108. (1) (NCERT XII, Pg 71, based on INHERITANCE OF ONE GENE)

109. (3) (NCERT XII, Pg 80, based on Law of Independent Assortment and Linkage)

110. (4) (NCERT XII, Pg 78, based on para 2)

111. (3) (NCERT XII, Pg 107, based on Figure 6.8)

112. (4) [NCERT class XI, Page 247, (Conceptual) Point 15.4.1 (Second paragraph)]

113. (2) [NCERT class XI exemplar, Question number 05]

114. (2) [NCERT Exemplar problems - XI page No. 22; Q.No.7]

115. (4) (NCERT XI Pg.227, Intro Part)

116. (3) (NCERT XI Pg.235, 14.6 Last line)

117. (3) (Exemplar 11<sup>th</sup> SAT Questions.3)

118. (3) (NCERT 11<sup>th</sup>, Living World, Exemplar)

119. (3) (NCERT XII, Pg 84, based on Figure 5.11)

120. (2) (NCERT 12<sup>th</sup>, SRFP, Exemplar)

121. (1) (11<sup>th</sup> NCERT exercises Page no.141)

122. (1) (NCERT 12<sup>th</sup>, SRFP, Exemplar)

123. (2) (NCERT 12<sup>th</sup>, SRFP, Exemplar)

124. (4) (NCERT XI – conceptual – page no. 223, 13.10.2, 2<sup>nd</sup> paragraph)

125. (3) (NCERT XI – Page no. 216, 13.7.2, 1 and page no. 219, 2<sup>nd</sup> paragraph)

126. (2) [NCERT Exemplar problems - XI page No. 22; Q.No.9]

127. (4) (NCERT XII, Pg 89, Based on colourblindness (Sex linkage))

128. (4) (11<sup>th</sup> NCERT EXEMPLAR, page no. 40)

129. (2) (11<sup>th</sup> NCERT EXEMPLAR, page no.39)

130. (3) [NCERT exemplar Class XI, Question number 08]

131. (3) (Exemplar 11<sup>th</sup> Question No.5)

132. (3) (NCERT XII, Pg 120, Salient Features of Human Genome)

133. (4) (NCERT XII, Pg 99, based on Packaging of DNA Helix)

134. (4) (NCERT XII, 115, para 3, line 1)

135. (3) (Exemplar 11<sup>th</sup> SAT Questions.5th)

**SECTION - B (Attempt Any 10 Questions)**

136. (4) (11<sup>th</sup> PK NCERT Page no 44 Exercises question bank conceptual)
137. (4) (NCERT-XII, Pg. 118)
138. (1) (NCERT XI – page no. 212, fig. 13.5 and 1<sup>st</sup> paragraph, page no. 214, fig. 13.7)
139. (3) (NCERT 11<sup>th</sup>, Living World, Exemplar)
140. (4) (NCERT 11<sup>th</sup>, Biological Classification, Exemplar)
141. (4) (NCERT-XII, Pg. 115, Para 2, Last 2<sup>nd</sup> line)
142. (2) (Exemplar 12<sup>th</sup> Question No.06)
143. (3) (11<sup>th</sup> NCERT EXERCISES Page no.141)
144. (2) (11<sup>th</sup> NCERT EXEMPLAR / PAGE NO. 51)
145. (2) [NCERT Exemplar problems - XI page No. 23; Q.No.10]
146. (3) [NCERT Class XI, Page no. 92, Line no. 07-09 (Conceptual)]
147. (2) [NCERT exemplar Class XI,, Question number 01]
148. (2) (NCERT 12<sup>th</sup>, SRFP, Exercise)
149. (3) (11<sup>th</sup> NCERT EXEMPLAR/ PAGE NO. 50)
150. (4) (NCERT XI Pg.228, 14.2 Last Para, 6<sup>th</sup> line)

**ZOOLOGY****Section - A (35 Questions)**

151. (4) (NCERT Page No. 146, Common diseases)
152. (1) (NCERT Page No. 157, Cancer)
153. (3) (Exemplar 12<sup>th</sup> Question No.08)
154. (2) (Page No. 297; 2nd line of 3rd paragraph)
155. (3) [NCERT P.No.302, 1<sup>st</sup> para & 303, 2<sup>nd</sup> para Line]
156. (3) (12<sup>th</sup> NCERT EXEMPLAR / PAGE NO. 70)
157. (2) [NCERT P.No.310, Last Para]
158. (2) [NCERT P.No.196 ,Para just above diagram]
159. (2) [NCERT P.No.211 ,Gene therapy 2<sup>nd</sup> para 2<sup>nd</sup> line]
160. (3) (NCERT 12<sup>th</sup> exemplar)
161. (3) (NCERT P.No.287, Disorders)
162. (3) (Exemplar 12<sup>th</sup> Question No.09)
163. (2) [NCERT P.No.310, 4<sup>th</sup> Line]
164. (3) [NCERT P.No.321, Last Para]
165. (4) (NCERT 11<sup>th</sup> exemplar)
166. (2) (NCERT P.No.274, Transport of oxygen)
167. (1) (NCERT 12<sup>th</sup> summary, p.no 55)

168. (4) (NCERT 11<sup>th</sup> exemplar frog)
169. (1) (NCERT P.No.271, Respiratory volume)
170. (2) (NCERT 12<sup>th</sup>, Evolution, Exemplar)
171. (3) (NCERT 12<sup>th</sup> exemplar)
172. (1) (NCERT 11<sup>th</sup> page no 122, EXERCISE)
173. (4) [NCERT P.No.316, 21.1 AND 21.2]
174. (1) [NCERT P.No.319, Fig. 21.3]
175. (4) (NCERT XI Page No.50 Phylum ctenophora)
176. (1) (NCERT XI Page No. 338; 4th paragraph)
177. (1) (NCERT 11<sup>th</sup>, Biomolecule, Exemplar)
178. (4) (NCERT P.No. 189)
179. (2) (NCERT XI Page No. 54; Phylum-echinodermata)
180. (2) (NCERT 11<sup>th</sup> exemplar cockroach)
181. (1) (NCERT 11<sup>th</sup>, Biomolecule, Exemplar)
182. (1) (NCERT 12<sup>th</sup>, Evolution, Exercise based)
183. (4) (NCERT 12<sup>th</sup>, Evolution, Summary)
184. (4) (12<sup>th</sup> NCERT EXEMPLAR / PAGE NO. 70 )
185. (3) (12<sup>th</sup> NCERT ecology 1<sup>st</sup> chapter, exponential equation— $dN/dt=(b-d) \times N$  Rate of change in population=2015-2005=10 Years  $dN/10=(0.028-0.008) \times 14=0.28 \times 10=2.8$ =Initial population 14 million  
Total population =2.8 million +14 million=16.8 near about 17 million )

**SECTION - B (Attempt Any 10 Questions)**

186. (4) [NCERT P.No.209, Pest Resistant plants]
187. (4) [NCERT P.No.321, 1<sup>st</sup> para 20<sup>th</sup> Line]
188. (3) (NCERT XI Page No. 291; 5th line of 2nd paragraph)
189. (1) [NCERT P.No.307, 9<sup>th</sup> Line]
190. (1) (NCERT P.No. 273, Exchange of gases)
191. (2) (NCERT Page No. 138)
192. (1) (NCERT Page No. 155, AIDS)
193. (4) (NCERT 12<sup>th</sup> exemplar)
194. (3) (NCERT 12<sup>th</sup> Exemplar)
195. (2) (11<sup>th</sup> NCERT Exercises / PAGE NO. 189)]
196. (1) (NCERT 11<sup>th</sup>, Biomolecule, Exemplar)
197. (2) (Exemplar 12<sup>th</sup> Question No.17)
198. (3) (NCERT 12<sup>th</sup>, Evolution, Exemplar)
199. (2) (Page No. 60, 3rd line)
200. (4) (NCERT XI Page No. 331; 5th line of 2nd paragraph)