

## ANSWER KEY & SOLUTION KEY FINAL ROUND - 17 (PCB) Dt.27.04.2024

### PHYSICS

#### SECTION - A (35 Questions)

01. (3)  $B_1 = \frac{\mu_0 I}{2R}, B_2 = \frac{2\mu_0 I}{2R}$

$$B_R = \sqrt{B_1^2 + B_2^2} = \frac{\mu_0}{2R} \sqrt{I^2 + (2I)^2}$$

$$\frac{\mu_0}{2R} \times \sqrt{5} I = \frac{\sqrt{5} \mu_0 I}{2R}$$

02. (1) Given,  $\vec{v} = (3\hat{i} + 5\hat{j})$  m/s

$$\vec{B} = (6\hat{i} + 4\hat{j})T$$


Magnetic force,  $\vec{F} = q(\vec{v} \times \vec{B})$

$$= -e[(3\hat{i} + 5\hat{j}) \times (6\hat{i} + 4\hat{j})]$$

$$= -e[(-30\hat{k} + 12\hat{k})] = 18e\hat{k}N$$

Thus, force is along positive Z-axis.

03. (1) We know in case of parallel plates the charges distributed as shown in the figure.



04. (4) According to Faraday's law of electro-magnetic induction Induced emf,  $e = \frac{Ldi}{dt}$

$$50 = L \left( \frac{5-2}{0.1 \text{ sec}} \right)$$

$$\Rightarrow L = \frac{50 \times 0.1}{3} = \frac{5}{3} = 1.67 \text{ H}$$

05. (3) Work function of aluminium is 4.2 eV. The energy of two photons can not be added at the moment photons collide with electron all its energy will be dissipated or wasted as this energy is

not sufficient to knock it out. Hence emission of electron is not possible.

06. (1) Current flowing through the conductor.

$$\frac{4}{1} = \frac{nev_{d1} \pi(1)^2}{nev_{d2} \pi(2)^2} \text{ or } \frac{v_{d1}}{v_{d2}} = \frac{4 \times 4}{1} = \frac{16}{1}$$

07. (1) According to Stefan's law

$$E = \sigma T^4$$

Heat radiated per unit area in 1 hour (3600s) is  
 $= 5 \times 10^{-8} \times (3000)^4 \times 3600 = 1.5 \times 10^{10}$

08. (4) Total charge  $Q_1 + Q_2 = Q'_1 + Q'_2$   
 $= 12\mu C - 3\mu C = 9\mu C$

Two isolated conducting spheres  $S_1$  and  $S_2$  are two connected by a conducting wire.

$$\therefore V_1 = V_2 = \frac{KQ'_1}{2/3R} = \frac{KQ'_2}{R/3} = 12 - 3 = 9\mu C$$

$$Q'_1 = 2Q'_2 \Rightarrow 2Q'_2 + Q'_2 = 9\mu C$$

$$\therefore Q'_1 = 6\mu C \text{ and } Q'_2 = 3\mu C$$

09. (1) When hot water temperature ( $T$ ) and surrounding temperature ( $T_0$ ) readings are noted, and  $\log(T - T_0)$  is plotted versus time, we get a straight line having a negative slope; as a proof of newton's law of cooling.

$$\frac{dT}{dt} = -K\Delta T$$

$$\int_{\Delta T_{initial}}^{\Delta T_{final}} \frac{dT}{\Delta T} = -K \int_0^t dt \Rightarrow \ln \left[ \frac{T - T_0}{T_i - T_0} \right] = -Kt$$

$$\Rightarrow \ln(T - T_0) = \ln(T_i - T_0) - kt$$

So on comparing  $y = -mx + c$

So option (1) is correct.

10. (1) Mass of section BC  $m/L(L - y)$ .


$$\therefore \text{tension at B} = T = m/L(L - y)g$$

$$\therefore \text{elongation of element } dy \text{ at B}$$

$$= dx = (dy) \frac{T}{AY} = \frac{m}{L} (L-y)g \frac{dy}{AY}$$

Total elongation =

$$\int dx = \frac{mg}{LAY} \int_0^L (L-y) dy = \frac{mgL}{2YA}$$



11. (3)  $\bar{M}$  (mag × moment / volume) =  $\frac{NiA}{Al}$

$$= \frac{Ni}{l} = \frac{(500)15}{25 \times 10^{-2}} = 30000 \text{ Am}^{-1}$$

12. (2) Majority carries in an n-type semiconductor are electrons.

13. (4)  $F + f = ma$  ... (i)

$$FR - fR = \frac{mR^2}{2} \frac{a}{R} \text{ ... (ii)}$$

From equation (i) and (ii)

$$2F = \frac{3ma}{2} \text{ or } a = \frac{4F}{3m}$$

$$F + f = \frac{m4F}{3m}$$

$$F = \frac{4}{3}F - F = \frac{F}{3}$$

14. (4) Translational KE =  $\frac{1}{2}mv^2$

Rotational KE =

$$\frac{1}{2}I\omega^2 = \frac{1}{2}mR^2\omega^2 \text{ (}\because \text{ For a ring, } I = mR^2 \text{)}$$

$$\therefore K_R = \frac{1}{2}m(\omega R)^2 = \frac{1}{2}mv^2$$

So translational KE = rotational KE


$$\therefore \frac{K_T}{K_R} = 1:1$$

15. (4)  $n = n$  to  $n = 1$ , number of transition =


$$\frac{n(n-1)}{2} = 10$$

$$n^2 - n = 20$$

$$n = 5$$



17. (1)



The voltage drop across  $1 \text{ k}\Omega = V_z = 15\text{V}$   
The current through  $1 \text{ k}\Omega$  is :

$$I' = \frac{15\text{V}}{1 \times 10^3 \Omega} = 15 \times 10^{-3} \text{ A} = 15\text{mA}$$

The voltage drop across  $250 \Omega = 20 \text{ V} - 15 \text{ V} = 5\text{V}$

The current through  $250 \Omega$  is

$$I = \frac{5\text{V}}{250\Omega} = 0.02 \text{ A} = 20\text{mA}$$

The current through the Zener diode is :

$$I_z = I - I' = (20 - 15)\text{mA} = 5\text{mA}$$

18. (1) Let pressure outside be  $P_0$

$$\therefore P_1 \text{ (in smaller bubble)} = P_0 + \frac{2T}{r}$$

$$P_2 \text{ (in bigger bubble)} = P_0 + \frac{2T}{R} \text{ (} R > r \text{)}$$

$$\therefore P_1 > P_2$$

Hence air moves from smaller bubble to bigger bubble.

19. (1)  $G = 15\Omega, i_g = 4 \text{ mA}, i = 6 \text{ A}$

Required shunt,

$$S = \left( \frac{i_g}{i - i_g} \right) G = \left( \frac{4 \times 10^{-3}}{6 - 4 \times 10^{-3}} \right) \times 15$$

$$\frac{4 \times 10^{-3}}{5.996} \times 15$$

=  $10\text{m}\Omega$  (in parallel)

20. (1) Kirchoff's loop rule follows from conservation of energy.

21. (1)  $\frac{\Delta A}{A} = 2 \frac{\Delta r}{r}$  [As  $A = 4\pi r^2$ ]

$\frac{\Delta V}{V} = 3 \frac{\Delta r}{r}$

$\therefore \frac{\Delta V}{V} = \frac{3 \Delta A}{2 A} \Rightarrow \frac{\Delta V}{V} = \frac{3}{2} \alpha$

22. (3)  $\phi = at + b, \phi + \Delta\phi = a(t + \Delta t) + b$ . Subtraction gives  $\Delta\phi = a\Delta t$

Average induced emf =  $(\Delta\phi / \Delta t) = a$ . The average induced current is  $a/R$

23. (4)  $n_1\beta_1 = n_2\beta_2$

$n_1 \left( \frac{D\lambda_1}{d} \right) = n_2 \left( \frac{D\lambda_2}{d} \right)$

$n_2 = n_1 \left( \frac{\lambda_1}{\lambda_2} \right) \Rightarrow 62 \times \frac{5893}{4358} \approx 84$

24. (3)  $(KE)_{\max} = E - \phi$

$= 1.8 - 1.2$

$= 0.6 \text{ eV}$

i.e.,  $eV_0 = (KE)_{\max} = eV_0 = 0.6 \text{ eV}$

$\therefore$  Stopping potential  $V_0 = 0.6 \text{ V}$

25. (1) A is false and B is true

26. (1) Conceptual

27. (2) By Gauss Law, flux is only by inside charges.

28. (1) Ideal ammeter has zero resistance

29. (4)  $U = U_1 + U_2$

$= n_1 C_{v1} T + n_2 C_{v2} T$

$= 3 \times \frac{5}{2} RT + 5 \times \frac{3}{2} R \times T$

$= 15 RT$

30. (4) Diffraction effect can be observed in both sound as well as light waves


31. (3) Magnitude of average velocity is

$|\vec{v}_{av}| = \left| \frac{\text{displacement}}{\text{time}} \right|$

$= \frac{PQ}{t} = \frac{\sqrt{2}R}{t}$

Here t can be found from

$\theta = \frac{1}{2} \alpha t^2$  or  $t = \sqrt{\frac{2\theta}{\alpha}}$ , where  $\theta = \frac{\pi}{2}$



$= \sqrt{\frac{2 \times \frac{\pi}{2}}{\pi/4}} = 2s \quad \therefore$

$|\vec{v}_{av}| = \frac{\sqrt{2}R}{t} = \frac{\sqrt{2} \cdot \sqrt{2}}{2} = 1 \text{ m/s}$

32. (2)  $E = Rhe \left[ \frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$

E will be maximum for the transition for which

$\left[ \frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$  is

Maximum. Here  $n_2$  is the higher energy level.

Clearly,  $\left[ \frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$  is maximum for the third transition,

i.e.  $2 \rightarrow 1$ . I transition represents the absorption of energy.

33. (2) Induced electric field is non conservative in nature and of circular in shape.

34. (3)  $\frac{d}{dv} (VP^n) = 0$

$vnP^{n-1} \frac{dP}{dv} + P^n = 0$

$nVP^{n-1} \frac{dP}{dv} = -P^n \quad \frac{dP}{dv} = \frac{-P^n}{nVP^{n-1}} = \frac{-P}{nV}$

Bulk modulus =  $\frac{dp}{-dV/V} = -V \frac{dP}{dv} = \frac{P}{n}$

35. (3) Volume of first substance,  $V_1 = 1/2$

Volume of second substance,  $V_2 = 4/3$

$\therefore$  Relative density =

$\frac{1+4}{(1/2) + (4/3)} = \frac{30}{11} = 2.73$

**SECTION - B (Attempt Any 10 Questions)**

36. (1)  $\frac{A_1}{A_2} = \frac{\sqrt{\beta}}{1}$ ,

$\frac{a_{\max}}{a_{\min}} = \frac{A_1 + A_2}{A_1 - A_2}$

37. (1) When interfering sources have same frequency and their phase difference remains constant with time, interference is sustained ( stayed for a finite time interval).

If amplitudes are of nearby values then contrast will be more pronounced.

38. (3)  $Z = \sqrt{R^2 + \left(2\pi fL - \frac{1}{2\pi fC}\right)^2}$

From above equation at  $f=0 \Rightarrow z = \infty$

When  $f = \frac{1}{2\pi\sqrt{LC}}$  (resonate frequency)  
 $\Rightarrow Z = R$


For  $f > \frac{1}{2\pi\sqrt{LC}} \Rightarrow Z$  starts increasing.

i.e., for frequency  $0 - f_r$ ,  $Z$  decreases and for  $f_r$  to  $\infty$ ,  $Z$  increases. This is justified by graph (3)

39. (4)


Force of friction will balance the weight.

So  $\mu Ma \geq Mg$ ;  $\mu = \frac{g}{a}$



40. (1) The equivalent circuit is shown in figure

Thus,  $C_{ab} = C = \frac{\epsilon_0 A}{d}$



41. (1) Given  $T/2 = 0.5$  s

$\therefore T = 1$  s

Frequency,  $f = \frac{1}{T} = \frac{1}{1} = 1$  Hz

If  $A$  is the amplitude, then

$2A = 50$  cm  $\Rightarrow A = 25$  cm

42. (3) Both statement I and II are correct

43. (2)  $\frac{S_{4th}}{S_{3rd}} = \frac{u + \frac{1}{2}g(2t_1 - 1)}{u + \frac{1}{2}g(2t_2 - 1)}$

$20 + \frac{1}{2} \times 10(2 \times 4 - 1)$   
 $= \frac{20 + \frac{1}{2} \times 10(2 \times 3 - 1)}{20 + \frac{1}{2} \times 10(2 \times 4 - 1)}$

$= \frac{20 + 5(7)}{20 + 5(5)} = \frac{20 + 35}{20 + 25} = \frac{55}{45} \Rightarrow \frac{S_{3rd}}{S_{2nd}} = \frac{11}{9}$

44. (1)  $\frac{GmM}{R^n} = \frac{mv^2}{R} \Rightarrow \sqrt{\frac{M}{R^{n-1}}} = V$

$T = \frac{2\pi R}{v} = \frac{2\pi R}{\sqrt{GM}} R^{\left(\frac{n-1}{2}\right)} \Rightarrow T \propto R^{\frac{n+1}{2}}$

45. (4) For 16 g of helium,  $n_1 = \frac{16}{4} = 4$

For 16 g of oxygen,  $n_2 = \frac{16}{32} = \frac{1}{2}$

For mixture of gases,

$C_V = \frac{n_1 C_{V1} + n_2 C_{V2}}{n_1 + n_2}$  where  $C_V = \frac{f}{2} R$


$C_P = \frac{n_1 C_{P1} + n_2 C_{P2}}{n_1 + n_2}$  where  $C_P = \left(\frac{f}{2} + 1\right) R$

For helium,  $f = 3$ ,  $n_1 = 4$

For oxygen,  $f = 5$ ,  $n_2 = 1/2$

$\therefore \frac{C_P}{C_V} = \frac{\left(4 \times \frac{5}{2} R\right) + \left(\frac{1}{2} \times \frac{7}{2} R\right)}{\left(4 \times \frac{3}{2} R\right) + \left(\frac{1}{2} \times \frac{5}{2} R\right)} = \frac{47}{29} = 1.62$

46. (2)



We know that  $y = x \tan \theta \left(1 - \frac{x}{R}\right)$

Now, from figure


$\tan \alpha + \tan \beta = \frac{x \tan \theta \left(1 - \frac{x}{R}\right)}{x} + \frac{x \tan \theta \left(1 - \frac{x}{R}\right)}{(R-x)}$

On solving  $\tan \alpha + \tan \beta = \tan \theta$

47. (1) Conceptual

48. (3)  $i = \frac{\epsilon}{r + R}$

$$v = iR = \frac{\epsilon R}{r + R} \Rightarrow v = \frac{\epsilon}{1 + \frac{r}{R}}$$



49. (2) Heat given by water,

$$Q_1 = ms\Delta T = 200 \times 1 \times (25 - 10) = 3000 \text{ cal}$$

Heat absorbed by  $m$  gm of ice at  $-14^\circ\text{C}$  to convert into water at  $10^\circ\text{C}$  is :

$$Q_2 = (ms\Delta T)_{\text{ice}} + mL_{\text{ice}} + (ms\Delta T)_{\text{water}}$$

$$= m(0.5 \times 14 + 80 + 1 \times 10) = 97m$$

Hence,  $97m = 3000$  or  $m = 31 \text{ gm}$ .

50. (1)  $d \sin \theta = n\lambda$

for 3rd maxima  $n = 3$

$$\therefore \sin \theta = \frac{n\lambda}{d} = \frac{3 \times 589 \times 10^{-9}}{0.589}$$

or  $\theta = \sin^{-1}(3 \times 10^{-6})$

## CHEMISTRY


### SECTION - A (35 Questions)

51. (3)  $\text{HO}-\overset{\text{O}}{\parallel}{\text{P}}-\text{OH}$  it ionizes in three steps because

three  $-\text{OH}$  groups are present.

52. (1)  $\Delta E = 0$  for isothermal process.

53. (2)  $2XY \rightleftharpoons X_2 + Y_2$ ;  $K_c = 81$



$$K'_c = \sqrt{K_c} = \sqrt{81} = 9$$

54. (2) In nitrobenzene  $-\text{NO}_2$  is strong electron withdrawing group decreases the reactivity

55. (2) each double bonded Carbon must be connected to two different group

56. (2) The ligands with small value of  $\Delta_0$  are called weak field ligands whereas those with large value



of  $\Delta_0$  are called strong field ligands, hence  $\text{CN}^-$  causes more splitting than  $\text{H}_2\text{O}$  and  $\text{NH}_3$ .

57. (1) In  $\text{F}_2\text{O}$ , fluorine is more electronegative than oxygen and hence given oxidation number of  $-1$ .

58. (1) Conceptual

59. (4)  $\text{K}_2\text{Cr}_2\text{O}_7$ ,  $\text{KMnO}_4$  and  $\text{K}_2\text{CrO}_4$  are coloured due to charge transfer.

60. (1)  $\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$



We can see that one mole of  $\text{KMnO}_4$  accepts 5 electrons, whereas one mole of  $\text{Fe}(\text{C}_2\text{O}_4)$  loses 3 electrons.

$\therefore$  Number of moles of  $\text{KMnO}_4$  required to oxidise one mole of  $\text{Fe}(\text{C}_2\text{O}_4) = 3/5 = 0.6$  mole.

61. (1) Conceptual

62. (1) Lanthanoids are 14 elements in the VIth period (atomic number = 58 to 71) that are filling the 4f-sublevel.

63. (1) Molality,  $m = \frac{w_B}{m_B} \times \frac{1000}{w_A}$

$$b = \frac{c}{m_B} \times \frac{1000}{(a-c)}; m_B = \frac{c}{b} \times \frac{1000}{(a-c)}$$

64. (4) Rate of Reaction  $\propto$  stability of carbocation.

65. (2) Assertion is true but Reason is false. The correct form of Reason is :

In  $\text{NH}_3$ , bond angle reduces to  $107.5^\circ$  due to repulsion between lone pair on N and bond pairs between N and H.

66. (3) The given figure is showing positive deviation from Raoult's law.


$$P_A > P_A^\circ X_A$$

Thus A-B attractive force should be weaker than A-A and B-B attractive forces.


67. (2)  $\text{CH}_3\text{CHO} + \text{RMgX} \xrightarrow{\text{H}_2\text{O}} \text{CH}_3-\text{CH}(\text{OH})-\text{R}$

68. (4) Acetate

69. (3) The overall reaction of Daniell cell is



So, its cell representation will be as follows :





70. (3) Conceptual

71. (2) Mechanism of reaction as well as relative concentration of reactants decides that how many concentration terms affect the rate of reaction i.e., order of reaction.

72. (2)  $\text{O}_2^{2-}$

73. (1) Catalyst catalyse both forward and backward reaction by same extent, without changing  $\Delta G$  and  $K_c$ .

74. (2) 'A' is  $\text{CH} \equiv \text{CH}$ ; 'B' is  $\text{CH}_3\text{CHO}$ ; 'C' is  $\text{CH}_3-\text{CH}_3$ ; 'D' is  $\text{CH}_3\text{COOH}$



75. (4) Both Assertion and Reason are true and Reason is the correct explanation of Assertion



$\text{NaCl}$  dissociates in water and organic acids dimerises in benzene.

76. (2) A = B = pricric acid

77. (1) Hydrated size of ion  $\propto$  Charge density of ion  $\rightarrow \text{H}^+$


78. (3) (I)-(C), (II)-(D), (III)-(A), (IV)-(B)

79. (1) 4



81. (2) No. of atoms =  $N_A \times \text{No. of moles} \times 3$  (atomicity)  
 $= 6.023 \times 10^{23} \times 0.1 \times 3 = 1.806 \times 10^{23}$

82. (4) It is valid for single electron species only



4-Bromo-6-hydroxycyclohex-2-ene-1-carboxylic acid

84. (2)  $\text{H}_2\text{C}=\overset{1}{\text{C}}=\overset{2}{\text{C}}=\overset{3}{\text{CH}}-\overset{4}{\text{CH}_3}$

85. (2)  $\text{XeOF}_4$


### SECTION - B (Attempt Any 10 Questions)

86. (2)  $\ddot{\text{N}}\text{H}_2-\ddot{\text{N}}\text{H}_2$  Neutral ligand

It does not act as bidentate because when it acts as bidentate a three membered ring (chelate complex) will be formed, that will be highly strained.



87. (3)  $E_{\text{metal}} = \frac{W \times 96500}{It} = \frac{22.2 \times 96500}{2 \times 5 \times 60 \times 60} = 59.5$

Oxidation number of the metal =  $\frac{177}{59.5} = +3$




89. (2) A-II, B-I, C-IV, D-III

90. (3)  $\text{C}(\text{graphite}) \rightarrow \text{C}(\text{diamond}), \Delta H = 1.9 \text{ kJ}$




$$(-\Delta H_1) - (-\Delta H_2) = 1.9 \text{ kJ or } \Delta H_2 = \Delta H_1 + 1.9$$

For combustion of 6 g,  $\Delta H_2 > \Delta H_1$  by  $1.9/2 = 0.95 \text{ kJ}$



Complex ion	Hybridization of central atom
$[\text{Fe}(\text{CN})_6]^{4-}$	$d^2sp^3$ (inner)
$[\text{Mn}(\text{CN})_6]^{4-}$	$d^2sp^3$ (inner)
$[\text{Co}(\text{NH}_3)_6]^{3+}$	$d^2sp^3$ (inner)
$[\text{Ni}(\text{NH}_3)_6]^{2+}$	$sp^3d^2$ (outer)



Total moles at equilibrium

$$= a - x + b - 3x + 2x = a + b - 2x$$

$$p_{\text{N}_2} = \text{Moles fraction of } \text{N}_2 \times P = \left( \frac{a-x}{a+b-2x} \right) P$$

$$p_{\text{H}_2} = \left( \frac{b-3x}{a+b-2x} \right) P; p_{\text{NH}_3} = \left( \frac{2x}{a+b-2x} \right) P$$

$$K_p = \frac{p_{\text{NH}_3}^2}{p_{\text{N}_2} \times p_{\text{H}_2}^3} = \frac{\left( \frac{2x}{a+b-2x} \right)^2 P^2}{\left( \frac{a-x}{a+b-2x} \right) P \left( \frac{b-3x}{a+b-2x} \right)^3 P^3}$$

$$= \frac{4x^2(a+b-2x)^2}{(a-x)(b-3x)^3 P^2}$$

94. (2) (I) Ketones do not give positive Tollen's and Fehling's test.

(II) Aromatic aldehydes do not give positive Fehling's test

(III) HCHO does not give positive haloform test

$$95. \quad (1) \text{ Energy of photon, } E = h\nu = \frac{hc}{\lambda}$$

$$= \frac{(6.6 \times 10^{-34} \text{ J s})(3 \times 10^8 \text{ ms}^{-1})}{5000 \times 10^{-10} \text{ m}} = 3.96 \times 10^{-19} \text{ J}$$

$$= \frac{3.96 \times 10^{-19} \text{ J}}{1.6 \times 10^{-19} \text{ J/eV}} = 2.475 \text{ eV}$$

( $\therefore 1 \text{ eV} = 1.6 \times 10^{-19} \text{ J}$ )

Kinetic energy of the emitted photon =  $h\nu - h\nu_0$

$$= 2.475 - 2.20 = 0.275 \text{ eV}$$

$$= 0.275 \times 1.6 \times 10^{-19} \text{ J} = 4.4 \times 10^{-20} \text{ J}$$

96. (4) The correct match is

A-II, B-III, C-IV, D-I

$E_{\text{cell}}^{\circ}$  can be determined by using this formula.

$$E_{\text{cell}}^{\circ} = E_{\text{right}} - E_{\text{left}} = E_{\text{cathode}} - E_{\text{anode}}$$

97. (3)  - Antiaromatic

98. (3) Due to inert effect the stability of lower oxidation state gradually increases while stability of higher oxidation state gradually decreases down the group in elements of group 13<sup>th</sup> to 15<sup>th</sup>. So correct orders are :

$$(iii) \text{ Pb}^{2+} > \text{Pb}^{4+}, \text{ Bi}^{3+} > \text{Bi}^{5+}$$

$$(iv) \text{ Sn}^{2+} < \text{Pb}^{2+}, \text{ Sn}^{4+} > \text{Pb}^{4+}$$

99. (2)  $k = \frac{2.303}{t} \log \frac{V_{\infty}}{V_{\infty} - V_t}$  gives constant value of k.

Hence, it is 1<sup>st</sup> order reaction

100. (2) The correct match is as

A-III, B-I, C-IV, D-II.

## BOTANY

### Section - A (35 Questions)

101. (3) (NCERT 11<sup>th</sup> Pg.237, 2<sup>nd</sup> line)
102. (2) (NCERT 12<sup>th</sup> Pg 122, Para 2)
103. (4) (NCERT 12<sup>th</sup> Pg 76, based on concept of Incomplete Dominance & law of independent assortment)
104. (3) ( NCERT 11<sup>th</sup> Page no.34 to 35, conceptual.)
105. (3) (NCERT 12<sup>th</sup> Pg 101, Biochemical Characterisation of Transforming Principle )

106. (2) (NCERT 12<sup>th</sup> Pg 102, based on Figure 6.5 )

107. (2) (NCERT 11<sup>th</sup>, Page no- 26, 2<sup>nd</sup> Paragraph, Line no- 1-12)

108. (3) (NCERT 12<sup>th</sup> Pg 87, Para 3, Line 16)

109. (3) (NCERT 12<sup>th</sup> Pg. 78 , Para 1, Line 4)

110. (3) (NCERT 12<sup>th</sup> Pg 106, 6.4.2)

111. (1) (NCERT 12<sup>th</sup> Pg 75, Figure 5.5)

112. (3) [NCERT 11<sup>th</sup> Newly added family]

113. (1) (NCERT 11<sup>th</sup>, Page no- 27, 2<sup>nd</sup> Paragraph, Line no- 5-7)

114. (1) (NCERT 11<sup>th</sup>, Page no- 7, 2<sup>nd</sup> paragraph, line no- 21,22)

115. (1) (NCERT 12<sup>th</sup>, Page no- 28, 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> paragraph, conceptual)

116. (1) (NCERT 11<sup>th</sup> Para 10.1.1, Page no.163 )

117. (1) (NCERT 11<sup>th</sup> Para 8.5.8, Page no.137 )

118. (2) (NCERT 11<sup>th</sup> Para8.5.3.4 , Page no.134 )

119. (2) (NCERT 11<sup>th</sup> Page no. 216, 13.7.2, Page no. 218, 13.8 - CONCEPT BASED and Page no. 220 1<sup>st</sup> paragraph)

120. (3) (NCERT 11<sup>th</sup> Page no. 212 – 19<sup>th</sup> line and Page 13.6 -concept based)

121. (4) [NCERT 11<sup>th</sup>, Page 248, point 15.4.3.1]

122. (1) [NCERT 11<sup>th</sup>, Page 248, Point 15.4.3.1]

123. (2) (NCERT 12<sup>th</sup> Page no. 247 fig.14.3 conceptual)

124. (1) (NCERT 11<sup>th</sup> Pg.231, 14.4.1 Last para 2<sup>nd</sup> line)

125. (4) (NCERT 12<sup>th</sup> Page no- 34, 1<sup>st</sup> paragraph, Concept based)

126. (3) (NCERT 11<sup>th</sup> Page no.31 fig.3.1, 32 to 33, name of the plant is fucus and its life cycle diplontic so diploid is dominant, not gametophyte.)

127. (1) (NCERT 12<sup>th</sup> Page No 38, Last para)

128. (2) (NCERT 12<sup>th</sup> Pg 117, based on Figure 6.14)

129. (4) (NCERT 12<sup>th</sup> Pg 85, 5.4)

130. (3) (NCERT 11<sup>th</sup> Para 10.1.1, Page no.163 )

131. (4) (NCERT 11<sup>th</sup> Page no.29 last line)

132. (3) [ NCERT 11<sup>th</sup>, Page no. 93 (First paragraph) and Point no. 6.3.4]

133. (3) [NCERT 11<sup>th</sup> Page No. 67; Sub-topic 5.1.2]

134. (2) [NCERT 11<sup>th</sup> Page No. 70; Sub-topic 5.3]

135. (3) (NCERT 12<sup>th</sup> Pg 85 based on polygenic)

### SECTION - B (Attempt Any 10 Questions)

136. (2) (NCERT 11<sup>th</sup>, Page no- 23, Paragraph-2<sup>nd</sup>, Line no-9-19)

137. (1) NCERT 12<sup>th</sup>, Page no- 34, Paragraph- 2.3, Line no- 5-8

138. (4) (NCERT 11<sup>th</sup>, Page no- 11, Table-1.1)  
 139. (1) (NCERT 11<sup>th</sup> Para 8.5.8, Page no.137 )  
 140. (4) (NCERT 11<sup>th</sup> Page no.37 fig.3.3 (d), salvinia is heterosporous so produces male and female gametophyte i.e. dioecious gametophyte.)  
 141. (4) (NCERT 11<sup>th</sup> Page no- 24, Paragraph- 2.3.3, Line no- 9 and 10)  
 142. (3) (NCERT 12<sup>th</sup> Page No. 246 2nd Para, 1st Line)  
 143. (2) [NCERT 11<sup>th</sup> Page No. 79, 80 & Newly added family]  
 144. (1) (NCERT 11<sup>th</sup> Para 8.5.8, Page no.137 )  
 145. (2) (NCERT 12<sup>th</sup> Pg 105, The Experimental Proof; Pg 106, Para 3; Pg 107, Para 2)  
 146. (4) [NCERT 11<sup>th</sup> , Page 249, Point 15.4.3.2]  
 147. (1) [NCERT 11<sup>th</sup> , Page no. 88, Subpoint 6.2.1]  
 148. (3) (NCERT 11<sup>th</sup> Para 10.2, 10.4 conceptual based, Page no.165-170 )  
 149. (3) (NCERT 11<sup>th</sup> Pg.230, 14.3, 3rd Paragraph, 1st line)  
 150. (2) (NCERT 11<sup>th</sup> Page no. 218, 1<sup>st</sup> paragraph)

## ZOOLOGY

### Section - A (35 Questions)

151. (2) (NCERT 12<sup>th</sup> page no 44, para1)  
 152. (2) (NCERT 11<sup>th</sup> page no 112, para1, line 16)  
 153. (2) (NCERT 11<sup>th</sup> Page No. 53; examples of arthropoda)  
 154. (4) (NCERT 12<sup>th</sup> Para 10.2.1 Page no.182 )  
 155. (3) (NCERT 12<sup>th</sup> NCERT Page no.260, 1<sup>st</sup> para last line and fig.15.1)  
 156. (4) (NCERT 11<sup>th</sup> Page No.- 183 - Respiratory organs)  
 157. (3) (NCERT 12<sup>th</sup> page no.60, last para)  
 158. (3) (NCERT 12<sup>th</sup> page no.64, Para 2, line 2)  
 159. (3) (NCERT 11<sup>th</sup> NCERT - Page No.- 196 - Coagulation of blood)  
 160. (3) [NCERT 12<sup>th</sup> P.No.195 2<sup>nd</sup> and 3<sup>rd</sup> Line]  
 161. (1) (NCERT 12<sup>th</sup> Page No.- 134 Immunity)  
 162. (3) (NCERT 11<sup>th</sup> Page No. 198)  
 163. (4) (NCERT 12<sup>th</sup> Page no.260, last line of 1<sup>st</sup> and fig.15.1, based)  
 164. (2) (NCERT 11<sup>th</sup> Page No. 336; Last line)  
 165. (2) (NCERT 11<sup>th</sup> P.No.309, Fig.20.6)  
 166. (2) [NCERT 11<sup>th</sup> P.No.312, Disorders]

167. (1) [NCERT 11<sup>th</sup> P.No.321, Line 9<sup>th</sup> to 12<sup>th</sup> ]  
 168. (2) [NCERT 11<sup>th</sup> P.No.321, 20<sup>th</sup> Line]  
 169. (3) (NCERT 11<sup>th</sup> Page No. 299; 1st line)  
 170. (1) (NCERT 11<sup>th</sup> Mixed question)  
 171. (4) (NCERT 11<sup>th</sup> Page No. 338, 2nd paragraph)  
 172. (4) (NCERT 12<sup>th</sup> Page no- 139, Figure 7.10)  
 173. (2) (NCERT 11<sup>th</sup>, Page no- 156, Figure-9.6)  
 174. (4) (NCERT 12<sup>th</sup> page no-129, 1<sup>st</sup> paragraph, line no- 11-13)  
 175. (3) [NCERT 12<sup>th</sup> P.No.198, 2<sup>nd</sup> para 3<sup>rd</sup> Line]  
 176. (2) [NCERT 12<sup>th</sup> P.No.312, 208 Last para]  
 177. (1) (NCERT 12<sup>th</sup> Page No.- 155 - Common Diseases)  
 178. (3) (NCERT 11<sup>th</sup>, Page no- 147, 2<sup>nd</sup> paragraph, Line no- 1<sup>st</sup> line)  
 179. (4) (NCERT 12<sup>th</sup> Para 10.1, 10.2.2, 10.5 Page no.181,182,187 )  
 180. (3) (NCERT 11<sup>th</sup> based extra)  
 181. (2) (NCERT 12<sup>th</sup> page no-128, 1<sup>st</sup> paragraph, line no-1 and 2)  
 182. (4) [NCERT 11<sup>th</sup> P.No.310, Last Para]  
 183. (2) (NCERT 12<sup>th</sup> page no. 52, factual)  
 184. (3) (NCERT 12<sup>th</sup> page no.62, para1)  
 185. (2) (NCERT 12<sup>th</sup> page no 43, para1)

### SECTION - B (Attempt Any 10 Questions)

186. (4) (NCERT 12<sup>th</sup> Page No. 142-143)  
 187. (4) [NCERT 11<sup>th</sup> P.No.304, Last Para]  
 188. (4) [NCERT 11<sup>th</sup> P.No.321, Forebrain and Hindbrain para]  
 189. (3) (NCERT 12<sup>th</sup> Page no.264 to 265.)  
 190. (4) (NCERT 12<sup>th</sup> Page no.229 1<sup>st</sup> para, 6<sup>th</sup> LINE)  
 191. (3) (NCERT 11<sup>th</sup> Page No.- 199 - Cardiac cycle)  
 192. (1) (NCERT 12<sup>th</sup> Page No.- 156 - Cancer)  
 193. (3) (NCERT 11<sup>th</sup> page no.104, para2)  
 194. (2) (NCERT 11<sup>th</sup> page no 102, last para)  
 195. (2) (NCERT 11<sup>th</sup> Page No. 297; 4th line)  
 196. (4) (NCERT 12<sup>th</sup> page no-127, 3<sup>rd</sup> paragraph, line no- 34)  
 197. (2) (NCERT 11<sup>th</sup>, Page no- 158, Paragraph- 9.12.5)  
 198. (4) (NCERT 12<sup>th</sup> Para 10.4 Page no.185 )  
 199. (3) [NCERT 12<sup>th</sup> P.No.213, 2<sup>nd</sup> and 3<sup>rd</sup> para]  
 200. (3) (NCERT 11<sup>th</sup> Page No. 339; 8th line of 4th Paragraph)