

Since 1999



CAREER INSTITUTE PVT.LTD.



NEET FRESH 2023-24

Mark
720

Group
PCB

PCB EXAM - 61

Date : 30/12/2023
Time : 3:20 Hours

Answer Key Version - Q (NEET FRESH All Batches)

| Physics | | | | | Chemistry | | | | |
|---------------|--------|--------|---------------|--------|----------------|--------|--------|---------------|--------|
| Sec. A | 11. 2 | 22. 2 | 33. 1 | 43. 2 | Sec. A | 61. 4 | 72. 3 | 83. 4 | 93. 4 |
| 01. 4 | 12. 1 | 23. 4 | 34. 2 | 44. 4 | 51. 4 | 62. 3 | 73. 2 | 84. 3 | 94. 1 |
| 02. 4 | 13. 3 | 24. 1 | 35. 3 | 45. 4 | 52. 1 | 63. 2 | 74. 3 | 85. 3 | 95. 2 |
| 03. 2 | 14. 2 | 25. 2 | Sec. B | 46. 4 | 53. 1 | 64. 1 | 75. 3 | Sec. B | 96. 2 |
| 04. 4 | 15. 2 | 26. 4 | 36. 2 | 47. 1 | 54. 4 | 65. 3 | 76. 2 | 86. 3 | 97. 1 |
| 05. 4 | 16. 3 | 27. 4 | 37. 1 | 48. 4 | 55. 2 | 66. 1 | 77. 2 | 87. 3 | 98. 1 |
| 06. 4 | 17. 4 | 28. 2 | 38. 1 | 49. 4 | 56. 1 | 67. 1 | 78. 4 | 88. 4 | 99. 1 |
| 07. 3 | 18. 4 | 29. 2 | 39. 3 | 50. 1 | 57. 2 | 68. 3 | 79. 2 | 89. 2 | 100. 1 |
| 08. 3 | 19. 3 | 30. 2 | 40. 3 | | 58. 4 | 69. 2 | 80. 1 | 90. 2 | |
| 09. 1 | 20. 2 | 31. 3 | 41. 1 | | 59. 3 | 70. 2 | 81. 4 | 91. 4 | |
| 10. 2 | 21. 1 | 32. 3 | 42. 1 | | 60. 2 | 71. 3 | 82. 2 | 92. 3 | |
| Biology | | | | | | | | | |
| Part-I | 110. 1 | 121. 1 | 132. 2 | 142. 3 | Part-II | 160. 3 | 171. 2 | 182. 1 | 192. 3 |
| Sec.A | 111. 2 | 122. 2 | 133. 2 | 143. 1 | Sec.A | 161. 2 | 172. 4 | 183. 2 | 193. 3 |
| 101. 1 | 112. 4 | 123. 3 | 134. 3 | 144. 4 | 151. 1 | 162. 2 | 173. 3 | 184. 4 | 194. 2 |
| 102. 3 | 113. 2 | 124. 2 | 135. 3 | 145. 3 | 152. 1 | 163. 3 | 174. 2 | 185. 3 | 195. 3 |
| 103. 4 | 114. 2 | 125. 1 | Sec.B | 146. 3 | 153. 3 | 164. 1 | 175. 2 | Sec. B | 196. 1 |
| 104. 3 | 115. 4 | 126. 1 | 136. 2 | 147. 4 | 154. 3 | 165. 2 | 176. 2 | 186. 4 | 197. 1 |
| 105. 4 | 116. 2 | 127. 1 | 137. 2 | 148. 4 | 155. 3 | 166. 2 | 177. 1 | 187. 1 | 198. 1 |
| 106. 4 | 117. 3 | 128. 1 | 138. 2 | 149. 4 | 156. 4 | 167. 1 | 178. 4 | 188. 3 | 199. 1 |
| 107. 1 | 118. 2 | 129. 2 | 139. 2 | 150. 2 | 157. 4 | 168. 2 | 179. 3 | 189. 4 | 200. 2 |
| 108. 2 | 119. 4 | 130. 4 | 140. 3 | | 158. 3 | 169. 3 | 180. 4 | 190. 3 | |
| 109. 1 | 120. 1 | 131. 1 | 141. 3 | | 159. 1 | 170. 3 | 181. 1 | 191. 2 | |

PHYSICS

SECTION - A (35 Questions)

01. (4) In the given situation, projectile could be considered as rigid body before explosion and after explosion, as its fragments are considered as system of particles. Thus, the concept of COM is applicable to both.

Since, the explosion is due to internal forces, so the motion of COM after explosion will follow the same parabolic path as it would have followed if there was no explosion.

Thus, statement given in option (4) is correct, rest are incorrect.

02. (4) $E_{\text{sphere}} = \frac{1}{2} I_s \omega^2 = \frac{1}{2} \times \frac{2}{5} MR^2 \times \omega^2$

$E_{\text{Cylinder}} = \frac{1}{2} I_c (2\omega)^2 = \frac{1}{2} \times \frac{MR^2}{2} \times 4\omega^2$


$\frac{E_{\text{sphere}}}{E_{\text{cylinder}}} = \frac{1}{5}$

03. (2) According to the equation of motion of the centre of mass. $M\vec{a}_{CM} = \vec{F}_{ext}$.

If $\vec{F}_{ext} = 0$, $\vec{a}_{CM} = 0$

i.e., $\vec{v}_{CM} = \text{constant}$

i.e., if no external force acts on a system (or resultant external force acting on a system is zero) the velocity of its centre of mass remains constant (i.e., velocity of the centre of mass is unaffected by internal forces).



Here, $F_{net} = 0$ but Kinetic energy will increase.

04. (4)

05. (4) Change in momentum

$= m\vec{v}_2 - m\vec{v}_1 = -mv - mv = -2mv$

06. (4) Resolve the 90N, 80N and 70N force into x and y components. The line of action of 90N, 50N, and x-components of the 80N and 70N forces pass through the pivot point A, therefore they cause no rotation.

∴ The total torque about point A is

$= (80 \sin 30^\circ) \left(\frac{L}{2}\right) - (60) \left(\frac{L}{2}\right) + (70 \cos 60^\circ)(L)$


$= (80) \left(\frac{1}{2}\right) \left(\frac{3}{2}\right) - (60) \left(\frac{3}{2}\right) + (70) \left(\frac{1}{2}\right) (3)$

$= 75 Nm$

07. (3) According to conservation of angular momentum

$I_1\omega_1 = I_2\omega_2 \Rightarrow \frac{1}{2} MR^2 \omega = \left(\frac{1}{2} MR^2 + \frac{1}{2} \left(\frac{M}{4}\right) R^2\right) \omega_2$

∴ $\omega_2 = \frac{4}{5} \omega$



Here, $m = 0.25 \text{ kg}$, $u_1 = 3 \text{ m/s}$, $u_2 = -1 \text{ m/s}$

It is an inelastic collision.

According to conservation of momentum

$mu_1 + mu_2 = (m + m)u$

$\Rightarrow v = \frac{mu_1 + mu_2}{2m} = \frac{u_1 + u_2}{2} = \frac{3 - 1}{2} = 1 \text{ m/s}$

09. (1) Given, $r = 0.4 \text{ m}$, $\alpha = 8 \text{ rad/s}^2$, $m = 4 \text{ kg}$, $I = ?$

Torque, $\tau = I\alpha = mgr \Rightarrow 4 \times 10 \times 0.4 = I \times 8$

$\Rightarrow I = \frac{16}{8} = 2 \text{ kg} \cdot \text{m}^2$

10. (2) $KE = \frac{L^2}{2I} \Rightarrow KE \propto \frac{1}{I}$ for same L

11. (2) When target is very light and at rest then after head on elastic collision it moves with double speed of projectile i.e. the velocity of body of mass m will be 2v.


12. (1) $U_1 = mgh_1$ and $U_2 = mgh_2$

% energy lost = $\frac{U_1 - U_2}{U_1} \times 100$

$\frac{mgh_1 - mgh_2}{mgh_1} \times 100 = \left(\frac{h_1 - h_2}{h_1}\right) \times 100$

$= \frac{2 - 1.5}{2} \times 100 = 25\%$

13. (3) From the triangle BCD



$$CD^2 = BC^2 - BD^2 = a^2 - \left(\frac{a}{2}\right)^2$$

$$x^2 = \frac{3a^2}{4} \Rightarrow x = \frac{\sqrt{3}a}{2}$$

Moment of inertia of system along the side AB

$$I_{\text{system}} = I_1 + I_2 + I_3 = m \times (0)^2 + m \times (x)^2 + m \times (0)^2$$

$$= mx^2 = m \left(\frac{\sqrt{3}a}{2}\right)^2 = \frac{3ma^2}{4}$$

14. (2)

15. (2) Here, R = 2m, M = 100 kg, v = 200 cm s⁻¹
= 20 × 10⁻² ms⁻¹

Total kinetic of the loop = K_T + K_R

$$= \frac{1}{2}Mv^2 + \frac{1}{2}I\omega^2 \quad [\because \text{For a hoop, } I = MR^2]$$

$$= \frac{1}{2}Mv^2 + \frac{1}{2}MR^2\omega^2$$

$$= \frac{1}{2}Mv^2 + \frac{1}{2}Mv^2 \quad [\because v = R\omega]$$

$$= Mv^2$$

Work required to stop the hoop = Total kinetic energy of the hoop.

$$Mv^2 = (100\text{kg})(20 \times 10^{-2} \text{ ms}^{-1})^2 = 4\text{J}$$

16. (3) $2u + m \times O = \frac{2u}{3} + mv$

$$\frac{4u}{3} = mv \quad \dots(1)$$

$$e = 1 = \frac{V - \frac{u}{3}}{u - 0}$$

$$V = \frac{4u}{3} \quad \dots(2)$$

On solving (1) and (2)

$$\frac{4u}{3} = m \times \frac{4u}{3}$$

$$\Rightarrow m = 1 \text{ kg.}$$

17. (4) $L = \sqrt{2IE}$. If E are equal then

$$\frac{L_1}{L_2} = \sqrt{\frac{I_1}{I_2}} = \sqrt{\frac{I}{2I}} = \frac{1}{\sqrt{2}}$$

18. (4) M.I. decreases and angular velocity increases.

19. (3) Force of friction

= K.E. of translation + K.E. of rotation

$$= \frac{1}{2}Mv^2 + \frac{1}{2}I\omega^2$$

$$= \frac{1}{2}Mv^2 + \frac{1}{2}Mk^2 \frac{v^2}{R^2}$$

$$(\because I = Mk^2 \text{ and } v = \omega R)$$

$$= \frac{1}{2}Mv^2 \left(1 + \frac{k^2}{R^2}\right)$$

$$\frac{\text{K.E. of rotation}}{\text{Total K.E.}} = \frac{\frac{1}{2}Mk^2 \frac{v^2}{R^2}}{\frac{1}{2}Mv^2 \left(1 + \frac{k^2}{R^2}\right)}$$


$$= \frac{\frac{k^2}{R^2}}{1 + \frac{k^2}{R^2}} = \frac{k^2}{k^2 + R^2}$$

20. (2) Velocity interchange when mass are equal


So K.E. = mgL.

21. (1) Conceptual

22. (2) $I_{cm} = \frac{ML^2}{12}$ (about middle point)



$$\therefore I = I_{cm} + Mx^2 = \frac{ML^2}{12} + M\left(\frac{L}{6}\right)^2 = \frac{ML^2}{9}$$



Weight of the rod will produce the torque.

$$\tau = I\alpha \Rightarrow mg \times \frac{l}{2} = \frac{ml^2}{3} \times \alpha$$

$$\text{Angular acceleration } \alpha = \frac{3g}{2l}$$

24. (1) Angular momentum of system remains constant

$$I \propto \frac{1}{\omega} \Rightarrow \frac{I_2}{I_1} = \frac{\omega_1}{\omega_2} = \frac{20}{10} \Rightarrow I_2 = 2I_1 = 2I$$

25. (2) Rotational kinetic energy = $\frac{1}{2} I\omega^2 = 1500$

$$\Rightarrow \frac{1}{2} \times 1.2 \times \omega^2 = 1500$$

$$\Rightarrow \omega^2 = \frac{3000}{1.2} \Rightarrow \omega = 50 \text{ rad/s}$$

Initially the body was at rest and after t sec its angular velocity becomes 50 rad/s.

$$\omega = \omega_0 + \alpha t \Rightarrow 50 = 0 + 25 \times t$$

$$\Rightarrow t = 2 \text{ s.}$$

26. (4) $f = \frac{KE_t}{KE_t + KE_r} = \frac{\frac{1}{2}mv^2}{\frac{1}{2}mv^2 + \frac{1}{2}I\omega^2} = \frac{5}{7}$

$$\left[\because \omega = \frac{v}{R}, I = \frac{2}{5}mR^2 \right]$$

27. (4)

28. (2) $E = \frac{L^2}{2I} \therefore E \propto L^2 \Rightarrow \frac{E_2}{E_1} = \left(\frac{L_2}{L_1}\right)^2$

$$\frac{E_2}{E_1} = \left(\frac{L_1 + 200\% \text{ of } L_1}{L_1}\right)^2 = \left(\frac{L_1 + 2L_1}{L_1}\right)^2 = (3)^2$$

$$\Rightarrow E_2 = 9E_1$$

Increment in kinetic energy

$$\Delta E = E_2 - E_1 = 9E_1 - E_1$$

$$\Delta E = 8E_1 \quad \Delta E = 8E_1 \quad \therefore \frac{\Delta E}{E_1} = 8 \text{ or percentage}$$

increase = 800%.

29. (2) $a = \frac{g \sin \theta}{1 + \frac{K^2}{R^2}}$


For disc $\frac{K^2}{R^2} = \frac{1}{2} = 0.5$

For sphere $\frac{K^2}{R^2} = \frac{2}{5} = 0.4$

a (sphere) > a (disc)

\therefore sphere reaches first

30. (2)



$$X_{com} = \frac{2000 \times 30}{2000 + 1000} = \frac{2000 \times 30}{3000} = 20 \text{ m.}$$

32. (3) A-2 ; B-1 ; C-4 ; D-3

Rolling motion \rightarrow combination of translatory and rotatory motion

Rate of change of angular momentum \rightarrow torque

$$\frac{dL}{dt} = \tau$$

Moment of inertia of a hollow cylinder about axis = MR^2

Theorem of parallel axis

$$I = I_{com} + Mr^2$$

33. (1) $X_{cm} = \frac{m_1x_1 + m_2x_2 + m_3x_3}{m_1 + m_2 + m_3}$

$$X_{cm} = \frac{300(0) + 500(40) + 400(70)}{300 + 500 + 400}$$

$$X_{cm} = \frac{500 \times 40 + 400 \times 70}{1200} \Rightarrow X_{cm} = 40cm.$$

34. (2) $X = \frac{m_1x_1 + m_2x_2 + m_3x_3 + m_4x_4}{m_1 + m_2 + m_3 + m_4}$

$$X = \frac{0 + 40x_4}{100} \Rightarrow 3 = \frac{40x_4}{100} \Rightarrow x_4 = \frac{300}{40} = 7.5$$

Similarly $y_4 = 7.5$ and $z_4 = 7.5$.

35. (3) According to law of conservation of linear momentum both pieces should possess equal momentum after explosion. As their masses are equal therefore they will possess equal speed in opposite direction.

Section - B (Attempt Any 10 Questions)


36. (2) As there is no external torque about the axis of rotation on the system of platform and tortoise, angular momentum will remain unchanged. As the tortoise moves moment of inertia of system will first decrease and then increase.

Hence angular velocity will first increase and then decrease. Also variation will not be linear.

37. (1) Mass of smaller circle = M

$$I = \frac{9MR^2}{2} - \left(\frac{1}{2} M \left(\frac{R}{3} \right)^2 + M \left(\frac{2R}{3} \right)^2 \right)$$

$$= 4MR^2.$$



When the cylinder is about to be raised, reaction at A vanishes. Taking torque about P.

$$F(2R - h) \geq Wl$$

$$l = \sqrt{R^2 - (R - h)^2}$$

$$= \sqrt{R^2 - \left(R - \frac{R}{3} \right)^2}$$

$$\Rightarrow \sqrt{R^2 - \frac{4R^2}{9}} = \frac{\sqrt{5}R}{3}$$

$$2R - h = 2R - \frac{R}{3} = \frac{5R}{3}$$


$$F \times \frac{5R}{3} \geq W \times \frac{\sqrt{5}R}{3}$$

$$F \geq \frac{W}{\sqrt{5}} \Rightarrow F_{min} = \frac{W}{\sqrt{5}}$$

39. (3) $h = e^{2n}h_0$

here, $n = 2$, so $\frac{h}{2} = e^4h$

$$\Rightarrow e = \left(\frac{1}{2} \right)^{1/4}$$




There is no external force so centre of mass will not shift.

41. (1) Angular momentum can be conserved about lowest point.

$$mv_0r + \frac{2}{5}mr^2 \frac{v_0}{2r} = mv'r + \frac{2}{5}mr^2 \frac{v'}{r}$$

$$\frac{10mv_0r + 2mv_0r}{10} = \frac{7mv'r}{5}$$

$$v' = \frac{6v_0}{7}$$



Particle falls from height h then formula for height covered by it in n th rebound is given by

$$h_n = e^{2n}h$$

where e = coefficient of restitution, n = No. of rebound Total distance travelled by particle before rebounding has stopped

$$H = h + 2h_1 + 2h_2 + 2h_3 + 2h_4 + \dots$$

$$= h + 2he^2 + 2he^4 + 2he^6 + 2he^8 + \dots$$

$$= h + 2h(e^2 + e^4 + e^6 + e^8 + \dots)$$

$$= h + 2h \left[\frac{e^2}{1-e^2} \right] = h \left[1 + \frac{2e^2}{1-e^2} \right] = h \left(\frac{1+e^2}{1-e^2} \right)$$

43. (2) $m = 2\text{kg}$, $V_1 = 0$, $V_2 = ?$, $t = 5$, $J = \text{Area}$

$$m(V_2 - V_1) = \frac{1}{2} \times 5 \times 10$$

$$2(V_2 - 0) = 25$$

$$V_2 = 12.5 \text{ m/s.}$$

44. (4) Apply conservation of angular momentum about the hinge.

$$mvR = \frac{m}{2} R^2 \omega + m(R\omega)R$$

$$R\omega = \frac{2v}{3} = \frac{2 \times 5}{3} = \frac{10}{3} \text{ m/s.}$$

45. (4) Rotational kinetic energy is

$$K_R = \frac{1}{2} I \omega^2 = \frac{1}{2} M k^2 \left(\frac{v}{R} \right)^2$$

$$(\because I = M k^2 \text{ and } v = R\omega)$$

$$\frac{1}{2} M v^2 \left(\frac{k^2}{R^2} \right)$$

Translational kinetic energy is

$$K_T = \frac{1}{2} M v^2$$

As per question, $K_R = 40\% K_T$

$$\therefore \frac{1}{2} M v^2 \left(\frac{k^2}{R^2} \right) = 40\% \frac{1}{2} M v^2$$

$$\text{or } \frac{k^2}{R^2} = \frac{40}{100} = \frac{2}{5}$$

$$\text{For a solid sphere, } \frac{k^2}{R^2} = \frac{2}{5}$$

Hence, the body is solid ball.

46. (4) According to parallel axes theorem

$$I = \frac{2}{5} m R^2 + m x^2$$

Hence graph (4) correctly depicts I vs x .

47. (1) By conservation of momentum,


$$m v + M \times 0 = (m + M) V$$

$$\text{Velocity of composite block } V = \left(\frac{m}{m + M} \right) v$$

$$\text{K.E. of composite block} = \frac{1}{2} (M + m) V^2$$

$$= \frac{1}{2} (M + m) \left(\frac{m}{M + m} \right)^2 v^2 = \frac{1}{2} m v^2 \left(\frac{m}{M + m} \right)$$

48. (4) From law of conservation of mechanical energy



$$\left[0 - \left(\frac{1}{2} I \omega^2 + \frac{1}{2} m v^2 \right) \right] + \left(m g \times \frac{3v^2}{4g} \right) = 0$$

$$\Rightarrow \frac{1}{2} I \omega^2 = \frac{3}{4} m v^2 - \frac{1}{2} m v^2 = \frac{m v^2}{2} \left(\frac{3}{2} - 1 \right) = \frac{m v^2}{4}$$

As cylinder is rolling $\omega = \frac{v}{R}$

$$\text{or } \frac{1}{2} I \frac{v^2}{R^2} = \frac{m v^2}{4}$$

$$\text{or } I = \frac{1}{2} m R^2$$

Hence, object is disc.

49. (4) Conceptual.

50. (1) $a = \frac{(3m - m)}{3m + m} g = \frac{g}{2}$


$$\vec{a}_{\text{cm}} = \frac{3m\vec{a}_1 + m\vec{a}_2}{3m + m}$$


Both mass have same magnitude of acceleration but in opposite direction $\vec{a}_1 = -\vec{a}_2 = a$ (Let)

$$a_{\text{cm}} = \left(\frac{3m - m}{4m} \right) \times \frac{g}{2} = \frac{g}{4}$$



CHEMISTRY


SECTION - A (35 Questions)

51. (4)
n-propyl propanoate
52. (1)
Diethyl ketone and methyl propyl ketone are position isomers
53. (1)
Molecular symmetry
54. (4)
Both (1) and (2)
55. (2)
2
56. (1)
A – (p), B – (r), C – (q), D – (s)
57. (2)
A – (p), B – (p), C – (q)
58. (4)

59. (3)
Infinite
60. (2)
The compound contains longest chain of 5C - atoms and e of ene is retained as the suffix name starts with constant
61. (4)
–NO₂ group, being strong electron-withdrawing, disperses the –ve charge, hence stabilizes the concerned carbanion.
62. (3)
In aqueous phase the decreasing order of basicity as

$$\text{C}_2\text{H}_5-\overset{\text{H}}{\text{N}}-\text{C}_2\text{H}_5 > \text{C}_2\text{H}_5-\underset{\text{C}_2\text{H}_5}{\text{N}}-\text{C}_2\text{H}_5 > \text{C}_2\text{H}_5-\text{NH}_2$$
63. (2)


64. (1)

$$\overset{\cdot}{\text{R}}-\overset{\cdot}{\text{X}} \xrightarrow[\text{light}]{\text{heat or}} \overset{\cdot}{\text{R}} + \overset{\cdot}{\text{X}}$$
65. (3)
CN[–], RCH₂[–], ROH
66. 
67. (1)
Cyclohexylamine
68. (3)
C₇H₇Cl has 4 isomers

69. (2)
The compound is a derivative of benzoic acid. The positions of substituents attached to benzene nucleus are represented by number of C-atoms and not by ortho, meta and para.
70. (2)
(a) > (d) > (b) > (c)
Due to S.I.P. effect (ortho effect) ortho methyl aniline is less basic than aniline.
71. (3)
b > a > c
72. (3)
2-ethyl-3-methylpentanal
73. (2)
Stability order of different alkyl carbocations on the basis of hyperconjugation is : 3° > 2° > 1° > methyl
In t-butyl cation, the C-atom bearing the positive charge is attached to three methyl groups therefore it possess nine α -hydrogens. It will give maximum nine hyperconjugative structures leading to maximum stability.
74. (3)
a, b, d
75. (3)
sp³, sp², sp
76. (2)



Antiform of butane is most stable conformer because of least steric and torsional strain.

77. (2)

Electromeric effect is purely a temporary effect and is brought into play only at the requirement of attacking reagent, it vanishes out as soon as the attacking reagent is removed from reaction mixture.

78. (4)

All of these

79. (2)

-OH shows +R effect while $>C=O$ shows -R effect.

80. (1)

A - (q), B - (p), C - (r), D - (s)


81. (4)

CH_3OH , CH_3CH_2OH


82. (2)

7

83. (4)



84. (3)



85. (3)

(c) > (b) > (d) > (a)

Because of ortho effect O-Toluic acid is more acidic than benzoic acid.

SECTION - B (Attempt Any 10 Questions)

86. (3)

$Cl_2C=CH-CH_2-CH_2-CH_2$. It can't show geometrical isomerism due to unsymmetrical alkene.


87. (3)

All resonating structures should have same number of electron pairs

88. (4)

Has two dissimilar groups attached to both ends of double bond

89. (2)



90. (2)

3

91. (4)

Assertion is incorrect, reason is correct


92. (3)

1, 1

93. (4)

$IV > III > I > II$

94. (1)




95. (2)

96. (2)

3, 4, 6-trimethyl octane

97. (1)



98. (1)

If both assertion and reason are true and the reason is the correct explanation of the assertion.

99. (1)

3-methyl-1-hexen-5-yne

100. (1)

If both assertion and reason are true and the reason is the correct explanation of the assertion.