

Since 1999



CAREER INSTITUTE PVT.LTD.



NEET FRESH 2023-24

Mark 720	Group PCB	PCB EXAM - 63	Date : 05/01/2024 Time : 3:20 Hours
-------------	--------------	----------------------	--

Answer Key Version - R (NEET FRESH All Batches)

Physics					Chemistry				
Sec. A	11. 3	22. 1	33. 4	43. 4	Sec. A	61. 2	72. 3	83. 2	93. 1
01. 4	12. 1	23. 4	34. 4	44. 2	51. 1	62. 2	73. 1	84. 1	94. 3
02. 1	13. 3	24. 3	35. 3	45. 2	52. 4	63. 2	74. 2	85. 4	95. 4
03. 2	14. 3	25. 4	Sec. B	46. 1	53. 3	64. 3	75. 3	Sec. B	96. 2
04. 1	15. 2	26. 4	36. 2	47. 2	54. 1	65. 4	76. 4	86. 4	97. 1
05. 3	16. 3	27. 4	37. 4	48. 2	55. 3	66. 1	77. 2	87. 1	98. 2
06. 2	17. 4	28. 4	38. 4	49. 4	56. 3	67. 4	78. 2	88. 4	99. 3
07. 1	18. 4	29. 1	39. 2	50. 2	57. 1	68. 1	79. 2	89. 3	100. 4
08. 1	19. 1	30. 4	40. 3		58. 2	69. 4	80. 3	90. 1	
09. 3	20. 4	31. 2	41. 4		59. 1	70. 4	81. 2	91. 2	
10. 1	21. 3	32. 2	42. 2		60. 1	71. 3	82. 3	92. 2	
Biology									
Part-I	110. 2	121. 1	132. 1	142. 3	Part-II	160. 4	171. 1	182. 3	192. 1
Sec.A	111. 3	122. 3	133. 2	143. 4	Sec.A	161. 2	172. 2	183. 3	193. 4
101. 3	112. 3	123. 3	134. 4	144. 4	151. 4	162. 2	173. 4	184. 2	194. 4
102. 3	113. 3	124. 4	135. 4	145. 2	152. 4	163. 3	174. 3	185. 2	195. 1
103. 2	114. 2	125. 4	Sec.B	146. 4	153. 4	164. 1	175. 2	Sec. B	196. 1
104. 3	115. 4	126. 3	136. 3	147. 2	154. 2	165. 4	176. 3	186. 2	197. 1
105. 4	116. 4	127. 1	137. 2	148. 3	155. 4	166. 2	177. 3	187. 4	198. 2
106. 1	117. 2	128. 1	138. 2	149. 3	156. 2	167. 2	178. 4	188. 3	199. 3
107. 3	118. 1	129. 3	139. 2	150. 3	157. 2	168. 3	179. 2	189. 2	200. 3
108. 3	119. 3	130. 4	140. 1		158. 4	169. 2	180. 2	190. 4	
109. 1	120. 1	131. 1	141. 4		159. 4	170. 4	181. 2	191. 4	

PHYSICS

SECTION - A (35 Questions)

01. (4) Particle velocity (v_p) = $-v \times$ slope of the graph at that point

At point 1 : Slope of the curve is positive, hence particle velocity is negative or downward (\downarrow).

02. (1) It is required to calculate the time from extreme position.

Hence, in this case equation for displacement of particle can be written as

$$x = A \sin\left(\omega t + \frac{\pi}{2}\right) = A \cos \omega t.$$

$$\text{So } \frac{A}{2} = A \cos \omega t \Rightarrow t = \frac{T}{6}.$$

03. (2) No of loops $\propto \frac{1}{\text{frequency}}$

$$\frac{f_1}{f_2} = \sqrt{\frac{T_1}{T_2}} = \sqrt{\frac{m_1}{m_2}}$$

$$2 = \sqrt{\frac{M}{M^1}} \Rightarrow M^1 = \frac{M}{4}.$$

04. (1)

$$x = A \cos\left(\omega t + \frac{\pi}{4}\right) \& v = \frac{dx}{dt} = -A\omega \sin\left(\omega t + \frac{\pi}{4}\right)$$

For maximum speed,

$$\sin\left(\omega t + \frac{\pi}{4}\right) = 1 \Rightarrow \omega t + \frac{\pi}{4} = \frac{\pi}{2} \text{ or } \omega t = \frac{\pi}{2} - \frac{\pi}{4}$$

$$\Rightarrow t = \frac{\pi}{4\omega}.$$

05. (3) The stone executes SHM about centre of earth

with time period $T = 2\pi\sqrt{\frac{R}{g}}$; where R = Radius of earth.

06. (2) From the given equation amplitude $A = 0.04$ m

$$\text{Frequency} = \frac{\text{Co-efficient of } t}{2\pi} = \frac{\pi/5}{2\pi} = \frac{1}{10} \text{ Hz}$$

$$\text{Wavelength } \lambda = \frac{2\pi}{\text{Co-efficient of } x} = \frac{2\pi}{\pi/9} = 18 \text{ m}$$

Wave speed

$$v = \frac{\text{Co-efficient of } t}{\text{Co-efficient of } x} = \frac{\pi/5}{\pi/9} = 1.8 \text{ m/s}.$$

07. (1) Maximum acceleration = $a\omega^2 = a \times 4\pi^2 n^2$
 $= 0.01 \times 4 \times (\pi)^2 \times (60)^2 = 144\pi^2 \text{ m/sec}.$

08. (1) $F = -KX \Rightarrow dW = Fdx = -KXdX$

$$\text{So, } \int_0^W dW = \int_0^W -KXdW \Rightarrow W = U = -\frac{1}{2} KX^2.$$

09. (3) Suppose at displacement y from mean position potential energy = kinetic energy

$$\Rightarrow \frac{1}{2} m(A^2 - y^2)\omega^2 = \frac{1}{2} m\omega^2 y^2$$

$$\Rightarrow A^2 = 2y^2 \Rightarrow y = \frac{A}{\sqrt{2}}.$$

10. (1) In first overtone of organ pipe open at one end,

$$n_c = \frac{3v}{4l_c} \dots\dots\dots(i)$$

Third harmonic or second overtone of organ pipe

$$\text{open at both ends, } n_0 = \frac{3v}{2l_0} \dots\dots\dots(ii)$$

$$\text{Under resonance } n_c = n_0 \Rightarrow \frac{3v}{4l_c} = \frac{3v}{2l_0} \Rightarrow \frac{l_c}{l_0} = \frac{1}{2}.$$

11. (3) The effective acceleration in a lift descending

$$\text{with acceleration } \frac{g}{3} \text{ is } g_{\text{eff}} = g - \frac{g}{3} = \frac{2g}{3}.$$

$$\therefore T = 2\pi\sqrt{\frac{L}{g_{\text{eff}}}} = 2\pi\sqrt{\frac{L}{2g/3}} = 2\pi\sqrt{\frac{3L}{2g}}.$$

12. (1) $f = \frac{1}{T} = \frac{1}{0.04} = 25 \text{ Hz}.$

13. (3) The given equation representing a wave travelling along $-y$ direction (because '+' sign is given between t term and x term).

On comparing it with $x = A \sin(\omega t + ky)$, we get

$$k = \frac{2\pi}{\lambda} = 12.58 \Rightarrow \lambda = \frac{2 \times 3.14}{12.58} = 0.5 \text{ m}.$$

14. (3) $E = \frac{1}{2} m\omega^2 A^2$

$$E' = \frac{1}{2} m \omega^2 \left(\frac{A}{2} \right)^2$$

$$E' = \frac{E}{4}$$

15. (2) i. $v = \omega \sqrt{A^2 - x^2} \Rightarrow \frac{v^2}{\omega^2 A^2} = 1 - \frac{x^2}{A^2}$

$$\Rightarrow \frac{v^2}{\omega^2 A^2} + \frac{x^2}{A^2} = 1$$

v vs x is ellipse.

ii. $a = -\omega^2 x$, a vs x is straight line.

iii. $U = \frac{1}{2} m \omega^2 x^2$, U vs x is parabola.

iv. $x = A \sin \omega t$, x vs t is sinusoidal.

16. (3) First overtone of closed organ pipe $n_1 = \frac{3v}{4l_1}$

Third overtone of open organ pipe $n_2 = \frac{4v}{2l_2}$

$$n_1 = n_2 \text{ (Given)} \Rightarrow \frac{3v}{4l_1} = \frac{4v}{2l_2} \Rightarrow \frac{l_1}{l_2} = \frac{3}{8}$$

17. (4) Points B and F are in same phase.

18. (4) $y = 5 \sin(\pi t + 4\pi)$, comparing it with standard

equation $y = a \sin(\omega t + \phi) = a \sin\left(\frac{2\pi t}{T} + \phi\right)$

$$a = 5m \text{ and } \frac{2\pi t}{T} = \pi t \Rightarrow T = 2 \text{ sec.}$$

19. (1) At point 2, the acceleration of the particle is maximum, which is at the extreme position. At extreme position, the velocity of the particle will be zero.

20. (4) The given equation can be written as

$$y = 4 \sin\left(4\pi t - \frac{\pi x}{16}\right) \Rightarrow (v) = \frac{\text{Co-efficient of } t(\omega)}{\text{Co-efficient of } x(K)}$$

$$\Rightarrow v = \frac{4\pi}{\pi/16} = 64 \text{ cm/s along } +x \text{ direction.}$$

21. (3) Given, $x = 10 \sin\left(2t - \frac{\pi}{6}\right)$

$$A = 10 \text{ and } \omega = 2 \text{ Hz}$$

$$\therefore v = \omega \sqrt{A^2 - x^2} = 2 \sqrt{(10)^2 - (6)^2}$$

$$= 2 \sqrt{100 - 36} = 2 \times 8 = 16 \text{ ms}^{-1}$$

22. (1) K.E. and P.E. completes two vibration in a time during which S.H.M. completes one vibration. Thus frequency of P.E. or K.E. is double than that of S.H.M.

23. (4) The second pendulum placed in a space laboratory orbiting around the earth is in a weightlessness state.

Hence, $g = 0$ so $T = \infty$

24. (3) As, we know, in SHM

Maximum acceleration of the particle,

$$\alpha = A\omega^2$$

Maximum velocity, $\beta = A\omega \Rightarrow \omega = \frac{\alpha}{\beta}$

$$\Rightarrow T = \frac{2\pi}{\omega} = \frac{2\pi\beta}{\alpha} \quad \left[\because \omega = \frac{2\pi}{T} \right]$$

25. (4) $y = \frac{1}{\sqrt{a}} \sin \omega t \pm \frac{1}{\sqrt{b}} \sin\left(\omega t + \frac{\pi}{2}\right)$

Here phase difference = $\frac{\pi}{2}$

\therefore The resultant amplitude

$$= \sqrt{\left(\frac{1}{\sqrt{a}}\right)^2 + \left(\frac{1}{\sqrt{b}}\right)^2} = \sqrt{\frac{1}{a} + \frac{1}{b}} = \sqrt{\frac{a+b}{ab}}$$

26. (4) $y = A \sin(\omega t + \phi) = A \sin\left(\frac{2\pi}{T} t + \phi\right)$

$$\Rightarrow y = 0.5 \sin\left(\frac{2\pi}{0.4} t + \frac{\pi}{2}\right)$$

$$\Rightarrow y = 0.5 \sin\left(5\pi t + \frac{\pi}{2}\right) = 0.5 \cos 5\pi t.$$

27. (4) $V_p = -V_w \times \text{slope}$.

28. (4) Acceleration = $-\omega^2 y$. So $F = -m\omega^2 y$.

y is sinusoidal function.

So F will be also sinusoidal function with phase difference π .

29. (1) Spring constant

$$k = \frac{F}{x} = \frac{10}{5 \times 10^{-2}} = 200 \text{ N/m}$$

And for spring-mass system undergoing SHM

$$T = 2\pi\sqrt{\frac{m}{k}} = 2\pi\sqrt{\frac{2}{200}} = \frac{2\pi}{10} = 0.628 \text{ s}$$

($\because m = 2 \text{ kg}$ given)

30. (4) At time $\frac{T}{2}; v = 0 \therefore$ Total energy Potential energy.

31. (2) From the equation $y(x,t) = 2a \sin kx \cos \omega t$ the position of nodes (where amplitude is zero) are given by $\sin kx = 0$ or $kx = n\pi$ where $n = 0, 1, 2, 3 \dots$

32. (2) The two springs on left side having spring constant of $2k$ each are in series, equivalent constant is k . The two springs on right hand side of mass M are in parallel. Their effective spring constant is $(k + 2k) = 3k$.

Equivalent spring constants of value k and $3k$ are in parallel and their net value of spring constant of all the four springs is $(k + 3k) = 4k$.

$$\therefore \text{Frequency of mass is } n = \frac{1}{2\pi} \sqrt{\frac{4k}{M}}$$

33. (4) After reflection from a rigid surface we know crest is reflected as trough. So final wave equation will be $y = -A \sin(kx + \omega t)$

34. (4) $T = 2\pi\sqrt{\frac{l}{g}} \Rightarrow T \propto \frac{l}{\sqrt{g}}$

$$\Rightarrow \frac{\Delta T}{T} \times 100 = -\frac{1}{2} \left(\frac{\Delta g}{g} \right) \times 100 = -\frac{1}{2} (-2\%) = 1\%$$

35. (3) From given equation

$$\omega = \frac{2\pi}{T} 0.5\pi \Rightarrow T = 4 \text{ sec}$$

Time taken from mean position to the maximum displacement = $\frac{1}{4} T = 1 \text{ sec}$.

Section - B (Attempt Any 10 Questions)

36. (2) We use equation $T = 2\pi\sqrt{\frac{I}{mgh}}$ and the parallel-axis theorem $I = I_{cm} + mh^2$ where $h = d$.

For a solid disk of mass m , the rotational inertia about its centre is $I_{cm} = mR^2 / 2$.

$$\text{Therefore, } T = 2\pi\sqrt{\frac{\frac{mR^2}{2} + \frac{mR^2}{4}}{mg \frac{R}{2}}} = 2\pi\sqrt{\frac{3R}{2g}}$$

37. (4)

$$v = \sqrt{\frac{T}{\mu}} = \sqrt{\frac{T}{\frac{\pi d^2}{4} \rho}} \Rightarrow v = 2\sqrt{\frac{T}{\mu d^2 \rho}} \Rightarrow v \propto \frac{\sqrt{T}}{d}$$

$$\frac{v_A}{v_B} = \sqrt{\frac{T_A}{T_B}} \times \frac{d_B}{d_A} = \frac{1}{\sqrt{2}} \times \frac{d_B}{d_B/2}$$


$$\Rightarrow \frac{v_A}{v_B} = \frac{2}{\sqrt{2}} \therefore v_A : v_B = \sqrt{2} : 1$$

38. (4)

$$T = \mu v^2 = \mu \frac{\omega^2}{k^2} = 0.04 \frac{(2\pi/0.004)^2}{(2\pi/0.50)^2} = 6.25 \text{ N}$$

39. (2) After 2s, the each wave travels a distance = $2 \times 2 = 4 \text{ m}$.

The wave shape is shown in figure. Thus energy is purely kinetic.



40. (3) $y = A \sin(\omega t - kx)$

Particle velocity,

$$v_p = \frac{dy}{dt} = A\omega \cos(\omega t - kx)$$

$$\therefore v_{p \max} = A\omega$$

$$\text{wave velocity} = \frac{\omega}{k}$$


$$\text{As per question } A\omega = \frac{\omega}{k}$$

$$\text{i. e., } A = \frac{1}{k} \text{ But } k = \frac{2\pi}{\lambda}$$

$$\therefore \lambda = 2\pi A$$

41. (4) For nodes $\sin(0.314x) = 0 = n$
 $\Rightarrow 0.314x = K_2\pi = K \times 3.14$
 $x = 10n$ [$n = 0, 1, 2, 3, \dots$]
 \therefore nodes are at 0, 10, 20, 30 cm
 $2\pi f = 600\pi \Rightarrow f = 300 \text{ Hz}$
 length $l = \frac{3\lambda}{2} = 3(10) = 30 \text{ cm}$
42. (2) Amplitude of a damped oscillator
 $A = A_0 e^{-bt/2m}$
 Case 1 : When $t = 2\text{s}$, $A = \frac{A_0}{3}$
 $\therefore \frac{A_0}{3} = A_0 e^{-2t/2m} \Rightarrow \frac{1}{3} = e^{-b/m} \dots\dots\dots(i)$
 When $t = 6\text{s}$, $A = \frac{A_0}{n}$
 $\therefore \frac{A_0}{n} = A_0 e^{-6b/2m} \Rightarrow \frac{1}{n} = (e^{-b/m})^3 \dots\dots\dots(ii)$
 From eqs. (i) and (ii), we get
 $\Rightarrow \frac{1}{n} = \left(\frac{1}{3}\right)^3 \Rightarrow \therefore n = 3^3$
43. (4) Suppose $n_p =$ frequency of piano = ?
 $(n_p \propto \sqrt{T})$
 $n_f =$ frequency of tuning fork = 256 Hz
 $x =$ beat frequency = 5 bps, which is decreasing
 $(5 \rightarrow 2)$ after clanging the tension of piano wire
 Also, tension of piano wire is increasing, so $n_p \downarrow$.
 Hence $n_p \uparrow - n_f = x \downarrow \longrightarrow$ Wrong
 $n_f - n_p \uparrow = x \downarrow \longrightarrow$ Correct.
 $\Rightarrow n_p = n_f - x = 256 - 5 \text{ Hz}$.
44. (2) As mg produces extension x , hence $k \Rightarrow \frac{mg}{x}$
 $\therefore T = 2\pi\sqrt{\frac{M+m}{k}} = 2\pi\sqrt{\frac{(M+m)x}{mg}}$
45. (2) $n = \frac{1}{2l}\sqrt{\frac{T}{m}} \Rightarrow n_1 l_1 = n_2 l_2 = n_3 l_3 = k$
 $l_1 + l_2 + l_3 = l \Rightarrow \frac{k}{n_1} + \frac{k}{n_2} + \frac{k}{n_3} = \frac{k}{n}$
 $\Rightarrow \frac{1}{n} = \frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3} + \dots\dots$

46. (1) K.E. of a body undergoing SHM is given by,
 $K.E. = \frac{1}{2}ma^2\omega^2 \cos^2 \omega t$ and $T.E. = \frac{1}{2}ma^2\omega^2$
 Given $K.E. = 0.75 \text{ T.E.}$
 $\Rightarrow 0.75 = \cos^2 \omega t \Rightarrow \omega t = \frac{\pi}{6}$
 $\Rightarrow t = \frac{\pi}{6 \times \omega} \Rightarrow t = \frac{\pi \times 2}{6 \times 2\pi} \Rightarrow t = \frac{1}{6} \text{ s}$
47. (2) $n_A =$ Known frequency = 288 cps, $n_B = ?$
 $x = 4 \text{ bps}$, which is decreasing (from 4 to 2) after loading i.e. $x \downarrow$
 Unknown fork is loaded, so $n_B \downarrow$
 Hence $n_A - n_B \downarrow = x \downarrow \longrightarrow$ Wrong
 $n_B \downarrow - n_A \downarrow = x \downarrow \longrightarrow$ Correct.
 $\Rightarrow n_B = n_A + x = 288 + 4 = 292 \text{ Hz}$.



- Given equations
 $y = A_0 + A \sin \omega t + B \cos \omega t$
 Now assume $(y - A_0) = \gamma$
 $y - A_0 = A \sin \omega t + B \cos \omega t$
 $\gamma = A \sin \omega t + B \cos \omega t$
 $= \sqrt{A^2 + B^2} \sin(\omega t + \phi)$
 which is S.H.M.
 where $\cos \phi = \frac{A}{\sqrt{A^2 + B^2}}$
 and $\sin \phi = \frac{B}{\sqrt{A^2 + B^2}}$
49. (4)
50. (2) $y = e^{-(ax^2 + bt^2 + 2\sqrt{ab}xt)}$
 $= e^{-(\sqrt{ax} + \sqrt{bt})^2}$
 $\sqrt{ax} + \sqrt{bt} = kx + \omega t$
 $k = \sqrt{a}, \omega = \sqrt{b}$
 Wave velocity, $v = \frac{\omega}{k} = \sqrt{\frac{b}{a}}$.

CHEMISTRY

SECTION - A (35 Questions)

51. (1)

Factual

52. (4)

Factual

53. (3)

Effect of increase of temperature on equilibrium constant depends on the fact that whether the reaction is exothermic, or endothermic. If the reaction is exothermic, it is favoured by low temperature and if the reaction is endothermic, it is favoured by high temperature.

54. (1)

Melting of ice involve absorption of heat i.e Endothermic hence high temperature favour the process. Further for a given mass volume of water is less than ice thus high pressure favour the process. High pressure and high temperature convert ice into liquid.

55. (3)

$$r_1 = k[A]^2$$

$$r_2 = \frac{k[A]^2}{4}$$

$$\frac{r_1}{r_2} = 4$$

$$r_2 = \frac{r_1}{4}$$

56. (3)

$$t = \frac{2.303}{k} \log \frac{100}{1}$$

$$= \frac{2.303}{2.303 \log 2} \times 2 \times 30$$


199 min

57. (1)

(A) only

58. (2)

A graph plotted between $\log k$ vs $\frac{1}{T}$ for calculating activation energy is shown as



from Arrhenius equation

$$\log k = \log A - \frac{E_a}{2.303 RT}$$

59. (1)

$$\Delta n_g = 0$$

$$K_p = K_c (RT)^{\Delta n_g}$$

$$K_p = K_c$$


$$\frac{K_p}{K_c} = 1$$

60. (1)

$$\text{Unit of } K_c = (\text{mol/L})^{\Delta n_g}$$

$$\text{mol L}^{-1}$$

61. (2)




62. (2)

The curve Y shows the increase in concentration of products with time.

63. (2)

Products predominate over reactant

64. (3)



$$4-x \quad 2x$$

$$4-x + 2x = 6$$

$$x = 2$$

$$K_p = \frac{16}{2} = 8$$

65. (4)

A - (s), B - (r), C - (p), D - (q)

66. (1)

A - (q), B - (r), C - (p)

67. (4)
Factual
68. (1)
 $\Delta G = \Delta G^0 + RT \ln Q$
69. (4)
In the given options $-\frac{d[C]}{3 \cdot dt}$ will not represent the reaction rate. It should not have -ve sign as it is product. Since $\frac{1}{3} \frac{dC}{dt}$ show the rate of formation of product C which will be positive.
70. (4)
 $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$
71. (3)
 $t_{1/2} \propto \frac{1}{a^{n-1}}$; where, a : initial concentration,
n : order of reaction
72. (3)
 $\frac{\ln 2}{k}$
73. (1)

A	→	2B	+	C
t = 0		P _i		0
t = t		P _i - y		2y

Total pressure P_t = P_i - y + 2y + y = P_i + 2y
 $\frac{P_t - P_i}{2} = y$
 $K = \frac{2.303}{t} \log \frac{2P_i}{3P_i - P_t}$
74. (2)
Rate $r_1 = k[A]^m [B]^n$... (1)
 $r_2 = 8 = k[2A]^m [2B]^n$... (2)
 $r_3 = 2 = k[A]^m [2B]^n$... (3)
By eqs. (1) and (3),
 $\frac{r_1}{r_3} = \frac{1}{2} = \left(\frac{1}{2}\right)^n$
n = 1
By eqs. (2) and (3),
 $\frac{r_3}{r_2} = \frac{2}{8} = \left(\frac{1}{2}\right)^m$
m = 2
∴ $r = k[A]^2 [B]^1$
75. (3)
Bimolecular reactions
76. (4)

$$K_c = \frac{K_p}{(RT)^{\Delta n}}$$

$$= \frac{1.44 \times 10^{-5}}{(0.082 \times 773)^{-2}} \text{ (R in L. atm. K}^{-1} \text{ mole}^{-1}\text{)}$$
77. (2)
 $\frac{1}{64}$
78. (2)
 $r = k [O_3]^2 [O_2]^{-1}$
79. (2)
The order w.r.t. I₂ is zero because the rate is not dependent on the concentration of I₂.
80. (3)
Reaction (3) can be obtained by adding reactions (1) and (2) therefore K₃ = K₁ · K₂
Hence (3) is the correct answer.
81. (2)

$$K_1 = \frac{1}{K_2} = \frac{1}{(K_3)^2}$$
82. (3)
k = (mol lit⁻¹)¹⁻ⁿ time⁻¹. For given reaction n = 2.
∴ k = mol⁻¹ lit sec⁻¹.
83. (2)

$$k = \frac{A_0}{2 t_{1/2}}$$

$$K_p = \frac{16}{2} = 8$$
84. (1)
 $K_p = K_c (RT)^{\Delta n}$
85. (4)
A - (q), B - (p), C - (r)

SECTION - B (Attempt Any 10 Questions)

86. (4)

$$\log \frac{k_2}{k_1} = \frac{E_a}{2.303R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$$

$$\log \frac{0.06}{0.03} = \frac{E_a}{2.303R} \left[\frac{500 - 400}{400 \times 500} \right]$$

$$E_a = 11.53 \text{ kJ}$$

87. (1)

Factual

88. (4)

For a zero order reaction,

$$t_{1/2} \propto a_0 \text{ (initial concentration or initial pressure)}$$

$$(t_{1/2})_1 \propto P_1$$

$$(t_{1/2})_2 \propto P_2$$

$$\frac{(t_{1/2})_2}{(t_{1/2})_1} = \frac{P_2}{P_1}, \quad \frac{(t_{1/2})_2}{45} = \frac{16}{4}$$

$$(t_{1/2})_2 = \frac{16}{4} \times 45 = 180 \text{ min}$$

89. (3)

(ii) and (iii)


90. (1)

A - (r), B - (p), C - (s), D - (q)

91. (2)

$$5.82 \times 10^{-2} \text{ atm}$$

92. (2)



$$a = 2, x = 0.4, V = 2L$$

$$\therefore [\text{PCl}_5] = \frac{2(1-0.4)}{2} = 0.6 \text{ mol L}^{-1}$$

$$[\text{PCl}_3] = [\text{Cl}_2] = \frac{2 \times 0.4}{2} = 0.4 \text{ mol L}^{-1}$$

$$\therefore K_c = \frac{0.4 \times 0.4}{0.6} = 0.267$$

93. (1)


For given reaction x and y may or may not be equal to p and q respectively.

94. (3)


$$\begin{aligned} t_{1/4} &= \frac{2.303}{k} \log \frac{1}{3/4} = \frac{2.303}{k} \log \frac{4}{3} \\ &= \frac{2.303}{k} (\log 4 - \log 3) = \frac{2.303}{k} (2 \log 2 - \log 3) \\ &= \frac{2.303}{k} (2 \times 0.301 - 0.4771) = \frac{0.29}{k} \end{aligned}$$

95. (4)

Given



$$K_1 = \frac{[\text{A}^+][\text{B}^-]}{[\text{AB}]}$$



$$K_2 = \frac{[\text{AB}_2^-]}{[\text{AB}][\text{B}^-]}$$

Dividing K_1 and K_2 , we get

$$K = \frac{K_1}{K_2} = \frac{[\text{A}^+][\text{B}^-]^2}{[\text{AB}_2^-]}$$

$$\therefore \frac{[\text{A}^+]}{[\text{AB}_2^-]} = \frac{K}{[\text{B}^-]^2}$$

96. (2)

An equilibrium constant does not give any information about the rate at which the equilibrium is reached.

97. (1)

$$t = \frac{2.303}{k} \log \frac{[\text{R}]_0}{[\text{R}]}$$

$$\log \frac{[\text{R}]_0}{[\text{R}]} = t \times \frac{k}{2.303}$$

$$y = mx$$

$$\frac{k}{2.303} = 0.02$$

$$k = 4.6 \times 10^{-2} \text{ sec}^{-1}$$

98. (2)

$$\frac{50}{t} = \frac{\ln 100 / 20}{\ln 100 / 10} \Rightarrow t = 50 \frac{\ln 10}{\ln 5}$$

99. (3)

Equilibrium constant is temperature dependent having one unique value for a particular reaction represented by a balanced equation at a given temperature.

100. (4)

The equilibrium will remain unaffected in all the three cases