

Since 1999



NEET 2023-24

Mark 720	Group PCB	PRE FINAL ROUND - 04	Date : 23/03/2024 Time : 3 :20 Hours
-------------	--------------	-----------------------------	---

Answer Key Version - P (PCB NEET 2023-24)

Physics					Chemistry				
Sec.A	11. 2	22. 4	33. 1	43. 1	Sec.A	61. 3	72. 1	83. 1	93. 3
01. 1	12. 2	23. 1	34. 1	44. 3	51. 3	62. 2	73. 3	84. 4	94. 2
02. 3	13. 2	24. 4	35. 1	45. 3	52. 3	63. 1	74. 1	85. 2	95. 3
03. 4	14. 1	25. 2	Sec. B	46. 4	53. 1	64. 3	75. 2	Sec. B	96. 1
04. 3	15. 3	26. 3	36. 4	47. 4	54. 4	65. 1	76. 1	86. 3	97. 4
05. 2	16. 1	27. 1	37. 3	48. 1	55. 1	66. 2	77. 2	87. 1	98. 3
06. 2	17. 3	28. 4	38. 2	49. 3	56. 1	67. 2	78. 4	88. 3	99. 3
07. 3	18. 2	29. 3	39. 4	50. 1	57. 1	68. 2	79. 3	89. 4	100. 1
08. 2	19. 2	30. 1	40. 3		58. 1	69. 4	80. 2	90. 1	
09. 1	20. 1	31. 2	41. 2		59. 1	70. 1	81. 3	91. 4	
10. 2	21. 2	32. 2	42. 3		60. 3	71. 2	82. 2	92. 3	
Biology									
Part-I Sec.A	110. 3	121. 1	132. 2	142. 4	Part-II Sec.A	160. 4	171. 4	182. 1	192. 1
	111. 1	122. 3	133. 2	143. 3		161. 2	172. 4	183. 3	193. 3
101. 1	112. 1	123. 2	134. 1	144. 2	151. 2	162. 2	173. 3	184. 2	194. 4
102. 3	113. 1	124. 1	135. 2	145. 2	152. 2	163. 2	174. 4	185. 4	195. 1
103. 1	114. 1	125. 3	Sec.B	146. 3	153. 4	164. 4	175. 4	Sec. B	196. 2
104. 1	115. 1	126. 3	136. 2	147. 3	154. 4	165. 1	176. 4	186. 3	197. 2
105. 4	116. 4	127. 2	137. 4	148. 1	155. 4	166. 1	177. 4	187. 2	198. 2
106. 1	117. 1	128. 2	138. 4	149. 4	156. 4	167. 3	178. 4	188. 4	199. 2
107. 1	118. 2	129. 1	139. 1	150. 1	157. 2	168. 1	179. 4	189. 4	200. 1
108. 2	119. 3	130. 3	140. 4		158. 4	169. 1	180. 3	190. 1	
109. 3	120. 1	131. 4	141. 2		159. 3	170. 3	181. 4	191. 1	

PHYSICS


SECTION - A (35 Questions)

01. (1) In vacuum velocity of all EM waves are same but their wavelengths are different.
02. (3) At the time $t=0$, e is maximum and is equal to E , but current i is zero.
As the time passes, current through the circuit increases but induced emf decreases.
03. (4) Peak value of r.m.s. value means, current become $\frac{1}{\sqrt{2}}$ times.

$$\text{So from } i = i_0 \sin 100\pi t \Rightarrow \frac{1}{\sqrt{2}} \times i_0 = i_0 \sin 100\pi t$$

$$\Rightarrow \sin \frac{\pi}{4} = \sin 100\pi t \Rightarrow t = \frac{1}{400} \text{ sec} = 2.5 \times 10^{-3} \text{ sec} .$$

04. (3) If pole strength, magnetic moment and length of each part are m' , M' and L' respectively then



$$m' = \frac{m}{2}$$

$$L' = L$$

$$\Rightarrow M' = \frac{M}{2}$$

$$m' = m$$

$$L' = \frac{L}{2}$$

$$\Rightarrow M' = \frac{M}{2}$$

05. (2) At $t=0$, phase of the voltage is zero, while phase of the current is $-\frac{\pi}{2}$ i.e., voltage leads by $\frac{\pi}{2}$.
06. (2) Potential across the PN junction varies symmetrically linear, having P side negative and N side positive.
07. (3) β - rays are beams of fast electrons.
08. (2) The energy of a photon is given by $E = \frac{hc}{\lambda} \Rightarrow E \propto \frac{1}{\lambda}$. Therefore, the graph of E v/s λ is rectangular hyperbola.


09. (1) $v = \frac{C}{\lambda}$

$$\Rightarrow v_1 = \frac{3 \times 10^8}{1} = 3 \times 10^8 \text{ Hz} = 300 \text{ MHz}$$

$$\text{and } v_2 = \frac{3 \times 10^8}{10} = 3 \times 10^7 \text{ Hz} = 30 \text{ MHz}$$

10. (2) $\left(\frac{d\phi}{dt}\right)_{\text{In first case}} = e$

$$\left(\frac{d\phi}{dt}\right)_{\text{relative velocity } 2v} = 2 \left(\frac{d\phi}{dt}\right)_{I \text{ case}} = 2e$$



11. (2) Repelled due to induction of similar poles.
12. (2) For a diamagnetic substance χ is small, negative and independent of temperature.
13. (2) This is because, when frequency ν is increased, the capacitive reactance $X_C = \frac{1}{2\pi\nu C}$ decreases and hence the current through the bulb increases.
14. (1) The stopping potential for curves a and b is same.

$$\therefore f_a = f_b$$

Also saturation current is proportional to intensity $\therefore I_a < I_b$.

15. (3) \vec{E} and \vec{B} are mutually perpendicular to each other and are in phase i.e. they become zero and minimum at the same place and at the same time.
16. (1) $|dq| = \frac{d\phi}{R} = i dt = \text{Area under } i-t \text{ graph}$
 $\therefore d\phi = (\text{Area under } i-t \text{ graph}) R$
 $= \frac{1}{2} \times 4 \times 0.1 \times (10) = 2 \text{ Wb} .$
17. (3) Time difference
 $= \frac{T}{2\pi} \times \phi = \frac{(1/50)}{2\pi} \times \frac{\pi}{4} = \frac{1}{400} \text{ s} = 2.5 \text{ m} - \text{s}$
18. (2) $\cos \phi = \frac{R}{Z} = \frac{R}{\sqrt{R^2 + \omega^2 L^2}}$
 $= \frac{12}{\sqrt{(12)^2 + 4 \times \pi^2 \times (60)^2 \times (0.1)^2}}$
 $\Rightarrow \cos \phi = 0.30$
19. (2) $E_0 = CB_0$


$$B_0 = \frac{9}{3 \times 10^8} = 3 \times 10^{-8} T$$

20. (1) Angular momentum is integral multiple of $\frac{h}{2\pi}$

$$mvr = \frac{nh}{2\pi}$$

So momentum $mv = \frac{nh}{2\pi r}$

21. (2) Input Output




22. (4) $r_n \propto n^2$

$$\Rightarrow \frac{r_4}{r_1} = \left(\frac{4}{1}\right)^2 = \frac{16}{1}$$

$$\Rightarrow r_4 = 16r_1$$


$$\Rightarrow r_4 = 16 r_0$$

23. (1)



$$C = \overline{\overline{A} \cdot \overline{B}} = \overline{\overline{A+B}} = A+B \text{ (De Morgan's theorem)}$$


Hence, Output C is equivalent to OR gate.



$$C = \overline{AB} \cdot AB = \overline{AB} + AB = AB + AB = AB$$

In this case output C is equivalent to AND gate.

24. (4) If E is the energy radiated in transition, then $E_{R \rightarrow G} > E_{Q \rightarrow S} > E_{R \rightarrow S} > E_{Q \rightarrow R} > E_{P \rightarrow Q}$
For getting blue line energy radiated should be maximum $\left(E \propto \frac{1}{\lambda}\right)$. Hence (4) is the correct option.



$$0 = m \cdot 2v - 2mv$$

mass \propto volume \propto (radius)³

$$\frac{r_1^3}{r_2^3} = \frac{m}{2m} \Rightarrow \frac{r_1}{r_2} = \left(\frac{1}{2}\right)^{1/3}$$

26. (3) $r \propto (A)^{1/3}$
27. (1)
28. (4) The output D for the given combination

$$D = \overline{(A+B)} \cdot \overline{C} = \overline{(A+B)} + \overline{C}$$

If $A = B = C = 0$

then $D = \overline{(0+0)} + \overline{0} = \overline{0} + \overline{0} = 1 + 1 = 1$

If $A = B = 1, C = 0$


then $D = \overline{(1+1)} + \overline{0} = \overline{1} + \overline{0} = 0 + 1 = 1$

29. (3)
30. (1) $V_n \propto \frac{Z}{n}$
 $v_n n = \text{constant}$
31. (2) $\lambda = \frac{h}{\sqrt{2mqV}} \Rightarrow \lambda \sqrt{V} = \text{constant}$

A rectangular hyperbola.

32. (2) Poles always occur in pair.

33. (1)



$$R = 1 \Omega$$

$$B = 0.5 T$$

$$\oint_B \vec{B} \cdot d\vec{s} = 0.5 \times 1 \times 1 \times \cos 0 = 0.5$$

34. (1) $\phi = 5t^3 + 4t^2 + 2t - 5$

$$|e| = \frac{d\phi}{dt} = 15t^2 + 8t + 2$$

At $t = 2, |e| = 15 \times 2^2 + 8 \times 2 + 2$

$$\Rightarrow e = 78V \Rightarrow I = \frac{e}{R} = \frac{78}{5} = 15.60$$


35. (1) $C = \frac{\omega}{k} = \frac{E_0}{B_0}$

SECTION - B (Attempt Any 10 Questions)

36. (4) From figure

$$\sin \frac{\theta}{2} = \frac{x}{r} \Rightarrow x = r \sin \frac{\theta}{2}$$

Hence new magnetic moment



- $$M' = m(2x) = m \cdot 2r \sin \frac{\theta}{2}$$
- $$= m \cdot \frac{2l}{\theta} \sin \frac{\theta}{2} = \frac{2ml \sin \theta / 2}{\theta} = \frac{2M \sin(\pi/6)}{\pi/3} = \frac{3M}{\pi}$$
37. (3) When magnet of length l is cut into four equal parts. then $m' = \frac{m}{2}$ and $l' = \frac{l}{2}$;
- $$\therefore M' = \frac{m}{2} \times \frac{l}{2} = \frac{ml}{4} = \frac{M}{4}$$
- New moment of inertia $I' = \frac{wl^2}{12}$
- $$= \frac{\frac{w}{4} \left(\frac{l}{2}\right)^2}{12} = \frac{1}{16} \cdot \frac{wl^2}{12}$$
- Here w is the mass of magnet
- $$\therefore I' = \frac{1}{16} I; \text{ Time period of each part}$$
- $$T' = 2\pi \sqrt{\frac{I}{M' B_H}}$$
- $$= 2\pi \sqrt{\frac{l/16}{(M/4) B_H}} = 2\pi \sqrt{\frac{l}{4MB_H}} = \frac{T}{2}$$
38. (2) $e = \frac{NBA(\cos \theta_2 - \cos \theta_1)}{\Delta t}$
- $$= -2000 \times 0.3 \times 70 \times 10^{-4} \frac{(\cos 180 - \cos 0)}{0.1}$$
- $$\Rightarrow e = 84 V$$
39. (4) $N\phi = Li \Rightarrow \frac{Nd\phi}{dt} = \frac{Ldi}{dt} \Rightarrow NB \frac{dA}{dt} = \frac{Ldi}{dt}$
- $$\Rightarrow \frac{1 \times 1 \times 5}{10^{-3}} = L \times \left(\frac{2-1}{2 \times 10^{-3}}\right) \Rightarrow L = 10 H$$
40. (3) $E = W_0 + K_{\max} \Rightarrow \frac{hc}{\lambda_1} = W_0 + E_1$
- and $\frac{hc}{\lambda_2} = W_0 + E_2$
- $$\Rightarrow hc = W_0 \lambda_1 + E_1 \lambda_1 \text{ and } hc = W_0 \lambda_2 + E_2 \lambda_2$$
- $$\Rightarrow W_0 \lambda_1 + E_1 \lambda_1 = W_0 \lambda_2 + E_2 \lambda_2$$
- $$\Rightarrow W_0 = \frac{E_1 \lambda_1 - E_2 \lambda_2}{(\lambda_2 - \lambda_1)}$$
41. (2) By using $E = W_0 + K_{\max} \Rightarrow K_{\max} = E - W_0$
- Hence, $K_1 = 1 - 0.5 = 0.5$ and $K_2 = 2.5 - 0.5 = 2$

$$\Rightarrow \frac{K_1}{K_2} = \frac{1}{4}.$$

42. (3) Since $W_0 = \frac{hc}{\lambda} - W_0$ and $2E = \frac{hc}{\lambda'} - W_0$

$$\lambda_T = \frac{\lambda_{Na} \times (W_0)_{Na}}{(W_0)_T} = \frac{5460 \times 2.3}{4.5} = 2791 \text{ \AA}$$

43. (1) For Lyman series

$$v_{Lyman} = \frac{c}{\lambda_{\max}} = Rc \left[\frac{1}{(1)^2} - \frac{1}{(2)^2} \right] = \frac{3RC}{4}$$

For Balmer series

$$v_{Balmer} = \frac{c}{\lambda_{\max}} = Rc \left[\frac{1}{(2)^2} - \frac{1}{(3)^2} \right] = \frac{5RC}{36}$$

$$\therefore \frac{v_{Lyman}}{v_{Balmer}} = \frac{27}{5}$$

44. (3) The Hydrogen atom before the transition was at rest. Therefore from conservation of momentum.

$$p_{H\text{-atom}} = p_{\text{photon}} = \frac{E_{\text{radiated}}}{c} = \frac{13.6 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) eV}{c}$$

$$1.6 \times 10^{-27} \times v = \frac{13.6 \left(\frac{1}{1^2} - \frac{1}{5^2} \right) \times 1.6 \times 10^{-19}}{3 \times 10^8}$$

$$\Rightarrow v = 4.352 \text{ m/s} \approx 4 \text{ m/sec}$$

45. (3) Energy is released in a process when total Binding energy (B.E.) of the nuclues is increased or we can say when total B.E. of products is more than the reactants. By calculation we can see that only in case of option (3), this happens.

Given $W \rightarrow 2Y$


$$\text{B.E. of reactants} = 120 \times 75 = 900 \text{ MeV}$$

$$\text{and B.E. of products} = 2 \times (60 \times 85)$$

$$= 1020 \text{ MeV}$$

i.e. B.E. of products > B.E. of reactants

46. (4) The electric field strength versus distance curve across the P-N junction is as follows



47. (4)

48. (1) $v_n \propto \frac{Z}{n}$, v_n v/s Z , inclined straight line

$$r_n \propto \frac{n^2}{Z}, r_n \text{ v/s } Z, \text{ rectangular hyperbola}$$

$$L_n = \frac{nh}{2\pi}, L_n \propto Z^0, \text{ parallel to x-axis, st. line}$$

$$E_n = -13.6 \frac{Z^2}{n^2}, \text{ rectangular hyperbola, } E_n \text{ is -ve}$$

49. (3) $\omega = 2\pi\nu = 2\pi \times 50 = 100\pi$

$$L = \frac{50}{\pi} \times 10^{-3} \text{ Hz}$$

$$C = \frac{10^3}{\pi} \times 10^{-6} = \frac{10^{-3}}{\pi}$$

$$R = 10 \Omega$$

$$X_C = \frac{1}{\omega C} = \frac{1}{100\pi \times \frac{10^{-3}}{\pi}} = 10$$

$$X_L = \omega L = 100\pi \times \frac{50}{\pi} \times 10^{-3} = 5.$$

$$\therefore Z = \sqrt{R^2 + (X_C - X_L)^2}$$

$$= \sqrt{10^2 + (10 - 5)^2} = \sqrt{100 + 25} = \sqrt{125} = 5\sqrt{5}.$$

50. (1) $\frac{n}{t} \times \frac{hc}{\lambda} = P$

$$\frac{n}{t} \times \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{600 \times 10^{-9}} = 3.3 \times 10^{-3} \frac{n}{t} = 10^{16}$$

CHEMISTRY


SECTION - A (35 Questions)

51. (3) Pentan-3-one can not show iodoform test here as it don't have $\text{CH}_3\text{CO-}$ gp.



52. (3) Etard's reaction

53. (1) Benzaldehyde

54. (4) $\Delta E = 0$, in a cyclic process.



56. (1) CCl_4 does not undergo hydrolysis at room temperature. Because C-atom does not have vacant orbital for accept lone pair electrons of H_2O molecule (Nucleophile)



58. (1) α -Keratin

59. (1) 5-Methyluracil

60. (3) Its dissociation constant is less as compound to carboxylic acids

61. (3) Hell-Volhard-Zelinsky reaction

62. (2) $100\pi\text{J}$

63. (1) As in process $\text{A} \rightarrow \text{B}$ volume is constant so it is isochoric.

In $\text{B} \rightarrow \text{C}$, pressure remains constant so it isobaric.


In $\text{C} \rightarrow \text{A}$, temperature remains constant so it isothermal.

64. (3) $\text{PH}_3 + \text{H}_3\text{PO}_4$

65. (1) Statement-I and Statement-II both are correct

66. (2) Lysine contains two basic groups. e.g., NH_2

67. (2) Retinol



69. (4) Statement-I is incorrect and Statement-II is correct.

70. (1) Number of equivalents of $\text{H}_2\text{C}_2\text{O}_4 = 2$

$$\text{Number of equivalents of } \text{K}_2\text{Cr}_2\text{O}_7 = \frac{1}{3} \times 6 = 2$$

71. (2) Peptization

72. (1) X-X bond F-F Cl-Cl Br-Br I-I
Bond dissociation energy (kcal/mol) 38 57 45.5 35.6

The lower value of bond dissociation energy of fluorine is due to the high inter-electronic repulsions between non-bonding electrons in the 2p-orbitals of fluorine. As a result F-F bond is weaker in comparison to Cl-Cl and Br-Br bonds.

73. (3) Due to larger size of iodine atom it can accommodate upto seven small fluorine atoms around, it while due to smaller sizes of chlorine and bromine atoms do not accommodate seven fluorine atoms, i.e., steric factor dominates in case of chlorine and bromine.

74. (1) Uracil

75. (2) Pyridine

76. (1) a-iv, b-ii, c-i, d-iii

77. (2) Sulphur only

78. (4) As for a pure substance T_A and T_B represent

