

Since 1999



# NEET 2023-24

Mark 720	Group PCB	<b>PRE FINAL ROUND - 04</b>	Date : 23/03/2024 Time : 3 :20 Hours
-------------	--------------	-----------------------------	---

## Answer Key Version - Q (PCB NEET 2023-24 )

Physics					Chemistry				
<b>Sec.A</b>	11. 1	22. 1	33. 4	43. 1	<b>Sec.A</b>	61. 2	72. 2	83. 1	93. 3
01. 2	12. 3	23. 4	34. 3	44. 3	51. 1	62. 2	73. 4	84. 4	94. 3
02. 3	13. 2	24. 3	35. 2	45. 1	52. 1	63. 2	74. 3	85. 1	95. 1
03. 2	14. 2	25. 1	<b>Sec. B</b>	46. 4	53. 1	64. 4	75. 2	<b>Sec. B</b>	96. 3
04. 1	15. 1	26. 2	36. 2	47. 3	54. 1	65. 1	76. 3	86. 4	97. 1
05. 2	16. 2	27. 2	37. 3	48. 2	55. 3	66. 2	77. 2	87. 3	98. 3
06. 2	17. 4	28. 1	38. 1	49. 4	56. 3	67. 1	78. 1	88. 3	99. 4
07. 2	18. 1	29. 1	39. 3	50. 3	57. 2	68. 3	79. 4	89. 2	100. 1
08. 2	19. 4	30. 1	40. 3		58. 1	69. 1	80. 2	90. 3	
09. 1	20. 2	31. 1	41. 4		59. 3	70. 2	81. 3	91. 1	
10. 3	21. 3	32. 3	42. 4		60. 1	71. 1	82. 3	92. 4	
Biology									
<b>Part-I Sec.A</b>	110. 1	121. 3	132. 3	142. 3	<b>Part-II Sec.A</b>	160. 1	171. 4	182. 2	192. 2
	111. 4	122. 2	133. 1	143. 1		161. 1	172. 4	183. 4	193. 2
101. 1	112. 1	123. 2	134. 1	144. 4	151. 4	162. 3	173. 4	184. 4	194. 2
102. 1	113. 2	124. 1	135. 4	145. 1	152. 2	163. 1	174. 4	185. 4	195. 1
103. 2	114. 3	125. 3	<b>Sec.B</b>	146. 2	153. 4	164. 1	175. 3	<b>Sec. B</b>	196. 3
104. 3	115. 1	126. 4	136. 2	147. 4	154. 3	165. 3	176. 4	186. 1	197. 2
105. 3	116. 1	127. 2	137. 4	148. 4	155. 4	166. 4	177. 1	187. 1	198. 4
106. 1	117. 3	128. 2	138. 3	149. 1	156. 2	167. 4	178. 3	188. 3	199. 4
107. 1	118. 2	129. 1	139. 2	150. 4	157. 2	168. 3	179. 2	189. 4	200. 1
108. 1	119. 1	130. 2	140. 2		158. 2	169. 4	180. 4	190. 1	
109. 1	120. 3	131. 1	141. 3		159. 4	170. 4	181. 2	191. 2	

# PHYSICS

## SECTION - A (35 Questions)

01. (2) Potential across the PN junction varies symmetrically linear, having P side negative and N side positive.

02. (3)  $\beta$  - rays are beams of fast electrons.

03. (2) The energy of a photon is given by  $E = \frac{hc}{\lambda} \Rightarrow E \propto \frac{1}{\lambda}$ . Therefore, the graph of E v/s  $\lambda$  is rectangular hyperbola.


04. (1)  $v = \frac{C}{\lambda}$

$$\Rightarrow v_1 = \frac{3 \times 10^8}{1} = 3 \times 10^8 \text{ Hz} = 300 \text{ MHz}$$

$$\text{and } v_2 = \frac{3 \times 10^8}{10} = 3 \times 10^7 \text{ Hz} = 30 \text{ MHz}$$

05. (2)  $\left(\frac{d\phi}{dt}\right)_{\text{In first case}} = e$

$$\left(\frac{d\phi}{dt}\right)_{\text{relative velocity } 2v} = 2 \left(\frac{d\phi}{dt}\right)_{\text{I case}} = 2e$$



06. (2) Repelled due to induction of similar poles.

07. (2) For a diamagnetic substance  $\chi$  is small, negative and independent of temperature.

08. (2) This is because, when frequency  $\nu$  is increased, the capacitive reactance  $X_C = \frac{1}{2\pi\nu C}$  decreases and hence the current through the bulb increases.

09. (1) The stopping potential for curves a and b is same.

$$\therefore f_a = f_b$$

Also saturation current is proportional to intensity

$$\therefore I_a < I_b.$$

10. (3)  $\vec{E}$  and  $\vec{B}$  are mutually perpendicular to each other and are in phase i.e. they become zero and minimum at the same place and at the same time.

11. (1)  $|dq| = \frac{d\phi}{R} = i dt = \text{Area under } i-t \text{ graph}$

$$\therefore d\phi = (\text{Area under } i-t \text{ graph}) R$$

$$= \frac{1}{2} \times 4 \times 0.1 \times (10) = 2 \text{ Wb}.$$

12. (3) Time difference

$$= \frac{T}{2\pi} \times \phi = \frac{(1/50)}{2\pi} \times \frac{\pi}{4} = \frac{1}{400} \text{ s} = 2.5 \text{ m-s}$$

13. (2)  $\cos \phi = \frac{R}{Z} = \frac{R}{\sqrt{R^2 + \omega^2 L^2}}$

$$= \frac{12}{\sqrt{(12)^2 + 4 \times \pi^2 \times (60)^2 \times (0.1)^2}}$$

$$\Rightarrow \cos \phi = 0.30$$

14. (2)  $E_0 = CB_0$

$$B_0 = \frac{9}{3 \times 10^8} = 3 \times 10^{-8} \text{ T}$$

15. (1) Angular momentum is integral multiple of  $\frac{h}{2\pi}$

$$mvr = \frac{nh}{2\pi}$$

$$\text{So momentum } mv = \frac{nh}{2\pi r}$$

16. (2) Input  $\rightarrow$  H.W.  $\rightarrow$  Output



Input, 60 Hz



Output, 60 Hz


17. (4)  $r_n \propto n^2$

$$\Rightarrow \frac{r_4}{r_1} = \left(\frac{4}{1}\right)^2 = \frac{16}{1}$$

$$\Rightarrow r_4 = 16r_1$$


$$\Rightarrow r_4 = 16r_0$$

18. (1)



$$C = \overline{A \cdot \overline{B}} = \overline{A} + B = A + B \text{ (De Morgan's theorem)}$$


Hence, Output C is equivalent to OR gate.



$$C = \overline{AB} \cdot \overline{AB} = \overline{AB + AB} = AB + AB = AB$$

In this case output C is equivalent to AND gate.

19. (4) If E is the energy radiated in transition, then  $E_{R \rightarrow G} > E_{Q \rightarrow S} > E_{R \rightarrow S} > E_{Q \rightarrow R} > E_{P \rightarrow Q}$ . For getting blue line energy radiated should be maximum  $\left(E \propto \frac{1}{\lambda}\right)$ . Hence (4) is the correct option.



$$0 = m \cdot 2v - 2mv$$

mass  $\propto$  volume  $\propto$  (radius)<sup>3</sup>

$$\frac{r_1^3}{r_2^3} = \frac{m}{2m} \Rightarrow \frac{r_1}{r_2} = \left(\frac{1}{2}\right)^{1/3}$$

21. (3)  $r \propto (A)^{1/3}$   
 22. (1)  
 23. (4) The output D for the given combination

$$D = \overline{(A+B)} \cdot \overline{C} = \overline{(A+B)} + \overline{C}$$

If A = B = C = 0

$$\text{then } D = \overline{(0+0)} + \overline{0} = \overline{0} + \overline{0} = 1 + 1 = 1$$


If A = B = 1, C = 0

$$\text{then } D = \overline{(1+1)} + \overline{0} = \overline{1} + \overline{0} = 0 + 1 = 1$$

24. (3)  
 25. (1)  $V_n \propto \frac{Z}{n}$   
 $v_n n = \text{constant}$   
 26. (2)  $\lambda = \frac{h}{\sqrt{2mqV}} \Rightarrow \lambda \sqrt{V} = \text{constant}$

A rectangular hyperbola.

27. (2) Poles always occur in pair.



$$R = 1 \Omega$$

$$B = 0.5 \text{ T}$$

$$\oint_B = \overline{B} \cdot \overline{ds} = 0.5 \times 1 \times 1 \times \cos 0 = 0.5$$

29. (1)  $\phi = 5t^3 + 4t^2 + 2t - 5$

$$|e| = \frac{d\phi}{dt} = 15t^2 + 8t + 2$$

$$\text{At } t = 2, |e| = 15 \times 2^2 + 8 \times 2 + 2$$


$$\Rightarrow e = 78V \Rightarrow I = \frac{e}{R} = \frac{78}{5} = 15.60$$

30. (1)  $C = \frac{\omega}{k} = \frac{E_0}{B_0}$   
 31. (1) In vacuum velocity of all EM waves are same but their wavelengths are different.  
 32. (3) At the time  $t = 0$ , e is maximum and is equal to E, but current i is zero. As the time passes, current through the circuit increases but induced emf decreases.  
 33. (4) Peak value of r.m.s. value means, current become  $\frac{1}{\sqrt{2}}$  times.

$$\text{So from } i = i_0 \sin 100\pi t \Rightarrow \frac{1}{\sqrt{2}} \times i_0 = i_0 \sin 100\pi t$$

$$\Rightarrow \sin \frac{\pi}{4} = \sin 100\pi t \Rightarrow t = \frac{1}{400} \text{ sec} = 2.5 \times 10^{-3} \text{ sec}$$

34. (3) If pole strength, magnetic moment and length of each part are  $m'$ ,  $M'$  and  $L'$  respectively then



$$m' = \frac{m}{2}$$

$$m' = m$$

$$L' = L$$

$$L' = \frac{L}{2}$$

$$\Rightarrow M' = \frac{M}{2}$$

$$\Rightarrow M' = \frac{M}{2}$$

35. (2) At  $t = 0$ , phase of the voltage is zero, while phase of the current is  $-\frac{\pi}{2}$  i.e., voltage leads by  $\frac{\pi}{2}$ .

**SECTION - B (Attempt Any 10 Questions)**

36. (2) By using  $E = W_0 + K_{\max} \Rightarrow K_{\max} = E - W_0$   
 Hence,  $K_1 = 1 - 0.5 = 0.5$  and  $K_2 = 2.5 - 0.5 = 2$   
 $\Rightarrow \frac{K_1}{K_2} = \frac{1}{4}$   
 37. (3) Since  $W_0 = \frac{hc}{\lambda} - W_0$  and  $2E = \frac{hc}{\lambda'} - W_0$

$$\lambda_T = \frac{\lambda_{Na} \times (W_0)_{Na}}{(W_0)_T} = \frac{5460 \times 2.3}{4.5} = 2791 \text{ \AA}$$

38. (1) For Lyman series

$$v_{Lyman} = \frac{c}{\lambda_{max}} = Rc \left[ \frac{1}{(1)^2} - \frac{1}{(2)^2} \right] = \frac{3RC}{4}$$

For Balmer series

$$v_{Balmer} = \frac{c}{\lambda_{max}} = Rc \left[ \frac{1}{(2)^2} - \frac{1}{(3)^2} \right] = \frac{5RC}{36}$$

$$\therefore \frac{v_{Lyman}}{v_{Balmer}} = \frac{27}{5}$$

39. (3) The Hydrogen atom before the transition was at rest. Therefore from conservation of momentum.

$$p_{H-atom} = p_{photon} = \frac{E_{radiated}}{c} = \frac{13.6 \left( \frac{1}{n_1^2} - \frac{1}{n_2^2} \right) eV}{c}$$

$$1.6 \times 10^{-27} \times v = \frac{13.6 \left( \frac{1}{1^2} - \frac{1}{5^2} \right) \times 1.6 \times 10^{-19}}{3 \times 10^8}$$

$$\Rightarrow v = 4.352 \text{ m/s} \approx 4 \text{ m/sec}$$

40. (3) Energy is released in a process when total Binding energy (B.E.) of the nuclues is increased or we can say when total B.E. of products is more than the reactants. By calculation we can see that only in case of option (3), this happens.

Given  $W \rightarrow 2Y$


$$\text{B.E. of reactants} = 120 \times 75 = 900 \text{ MeV}$$

$$\text{and B.E. of products} = 2 \times (60 \times 85)$$

$$= 1020 \text{ MeV}$$

i.e. B.E. of products > B.E. of reactants

41. (4) The electric field strength versus distance curve across the P-N junction is as follows



42. (4)

43. (1)  $v_n \propto \frac{Z}{n}$ ,  $v_n$  v/s  $Z$ , inclined straight line

$$r_n \propto \frac{n^2}{Z}, r_n \text{ v/s } Z, \text{ rectangular hyperbola}$$

$$L_n = \frac{nh}{2\pi}, L_n \propto Z^0, \text{ parallel to x-axis, st. line}$$

$$E_n = -13.6 \frac{Z^2}{n^2}, \text{ rectangular hyperbola, } E_n \text{ is -ve}$$

44. (3)  $\omega = 2\pi\nu = 2\pi \times 50 = 100\pi$

$$L = \frac{50}{\pi} \times 10^{-3} \text{ Hz}$$

$$C = \frac{10^3}{\pi} \times 10^{-6} = \frac{10^{-3}}{\pi}$$

$$R = 10 \Omega$$

$$X_C = \frac{1}{\omega C} = \frac{1}{100\pi \times \frac{10^{-3}}{\pi}} = 10$$

$$X_L = \omega L = 100\pi \times \frac{50}{\pi} \times 10^{-3} = 5$$

$$\therefore Z = \sqrt{R^2 + (X_C - X_L)^2}$$


$$= \sqrt{10^2 + (10 - 5)^2} = \sqrt{100 + 25} = \sqrt{125} = 5\sqrt{5}$$

45. (1)  $\frac{n}{t} \times \frac{hc}{\lambda} = P$

$$\frac{n}{t} \times \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{600 \times 10^{-9}} = 3.3 \times 10^{-9} \frac{n}{t} = 10^{16}$$

46. (4) From figure

$$\sin \frac{\theta}{2} = \frac{x}{r} \Rightarrow x = r \sin \frac{\theta}{2}$$



Hence new magnetic moment

$$M' = m(2x) = m \cdot 2r \sin \frac{\theta}{2}$$

$$= m \cdot \frac{2l}{\theta} \sin \frac{\theta}{2} = \frac{2ml \sin \theta / 2}{\theta} = \frac{2M \sin(\pi/6)}{\pi/3} = \frac{3M}{\pi}$$

47. (3) When magnet of length  $l$  is cut into four equal

parts. then  $m' = \frac{m}{2}$  and  $l' = \frac{l}{2}$ ;

$$\therefore M' = \frac{m}{2} \times \frac{l}{2} = \frac{ml}{4} = \frac{M}{4}$$

$$\text{New moment of inertia } I' = \frac{wl^2}{12}$$

$$= \frac{w}{12} \left( \frac{l}{2} \right)^2 = \frac{1}{16} \cdot \frac{wl^2}{12}$$

Here  $w$  is the mass of magnet

$$\therefore I' = \frac{1}{16} l; \text{ Time period of each part}$$

$$T' = 2\pi \sqrt{\frac{I'}{M' B_H}}$$

$$= 2\pi \sqrt{\frac{l/16}{(M/4)B_H}} = 2\pi \sqrt{\frac{l}{4MB_H}} = \frac{T}{2}$$

$$48. \quad (2) \quad e = \frac{NBA(\cos\theta_2 - \cos\theta_1)}{\Delta t}$$

$$= -2000 \times 0.3 \times 70 \times 10^{-4} \frac{(\cos 180 - \cos 0)}{0.1}$$

$$\Rightarrow e = 84 \text{ V}$$

$$49. \quad (4) \quad N\phi = Li$$

$$\Rightarrow \frac{Nd\phi}{dt} = \frac{Ldi}{dt} \Rightarrow NB \frac{dA}{dt} = \frac{Ldi}{dt}$$

$$\Rightarrow \frac{1 \times 1 \times 5}{10^{-3}} = L \times \left( \frac{2-1}{2 \times 10^{-3}} \right) \Rightarrow L = 10 \text{ H}$$

$$50. \quad (3) \quad E = W_0 + K_{\max}$$

$$\Rightarrow \frac{hc}{\lambda_1} = W_0 + E_1$$

$$\text{and } \frac{hc}{\lambda_2} = W_0 + E_2$$

$$\Rightarrow hc = W_0\lambda_1 + E_1\lambda_1 \quad \text{and} \quad hc = W_0\lambda_2 + E_2\lambda_2$$


$$\Rightarrow W_0\lambda_1 + E_1\lambda_1 = W_0\lambda_2 + E_2\lambda_2$$

$$\Rightarrow W_0 = \frac{E_1\lambda_1 - E_2\lambda_2}{(\lambda_2 - \lambda_1)}$$

## CHEMISTRY

### SECTION - A (35 Questions)


51. (1)  $\text{CCl}_4$  does not undergo hydrolysis at room temperature. Because C-atom does not have vacant orbital for accept lone pair electrons of  $\text{H}_2\text{O}$  molecule (Nucleophile)



52. (1)  ${}^{26}_{88}\text{Ra} \longrightarrow {}^{222}_{86}\text{Rn} + {}^4_2\text{He}$
53. (1)  $\alpha$ -Keratin
54. (1) 5-Methyluracil
55. (3) Its dissociation constant is less as compound to carboxylic acids
56. (3) Hell-Volhard Zelinsky reaction
57. (2)  $100\pi \text{ J}$
58. (1) As in process  $\text{A} \rightarrow \text{B}$  volume is constant so it is isochoric.  
In  $\text{B} \rightarrow \text{C}$ , pressure remains constant so it isobaric.

In  $\text{C} \rightarrow \text{A}$ , temperature remains constant so it isothermal.

59. (3)  $\text{PH}_3 + \text{H}_3\text{PO}_4$
60. (1) Statement-I and Statement-II both are correct
61. (2) Lysine contains two basic groups. e.g.,  $\text{NH}_2$
62. (2) Retinol



64. (4) Statement-I is incorrect and Statement-II is correct.

65. (1) Number of equivalents of  $\text{H}_2\text{C}_2\text{O}_4 = 2$

$$\text{Number of equivalents of } \text{K}_2\text{Cr}_2\text{O}_7 = \frac{1}{3} \times 6 = 2$$


66. (2) Peptization

67. (1) X-X bond      F-F    Cl-Cl    Br-Br    I-I  
Bond dissociation 38    57    45.5    35.6  
energy (kcal/mol)

The lower value of bond dissociation energy of fluorine is due to the high inter-electronic repulsions between non-bonding electrons in the 2p-orbitals of fluorine. As a result F-F bond is weaker in comparison to Cl-Cl and Br-Br bonds.


68. (3) Due to larger size of iodine atom it can accommodate upto seven small fluorine atoms around, it while due to smaller sizes of chlorine and bromine atoms do not accommodate seven fluorine atoms, i.e., steric factor dominate in case of chlorine and bromine.

69. (1) Uracil
70. (2) Pyridine
71. (1) a-iv, b-ii, c-i, d-iii
72. (2) Sulphur only
73. (4) As for a pure substance  $T_A$  and  $T_B$  represent the same temperature so, option 4 is correct here.
74. (3) A red coloured ppt. is obtained
75. (2) 1-iii, 2-iv, 3-i, 4-ii
76. (3) Statement-I is correct and Statement-II is incorrect
77. (2) Due to the absence of H-bonding,  $\text{PH}_3$  has the lowest b.p. The boiling point of the V group hydrides is :  
 $\text{BiH}_3 > \text{SbH}_3 > \text{NH}_3 > \text{AsH}_3 > \text{PH}_3$







Five chiral centres

79. (4) Abscisic acid (ABA)  
 80. (2)  $K_2Fe[Fe(CN)_6]$ —white]  
 81. (3) Pentan-3-one can not show iodoform test here as it don't have  $CH_3CO-$  gp.  
 82. (3) Etard's reaction  
 83. (1) Benzaldehyde  
 84. (4)  $\Delta E = 0$ , in a cyclic process.




### SECTION - B (Attempt Any 10 Questions)

86. (4) Tollen's reagent is not used in the detection of unsaturation but is used for distinction of (1) aldehydes from ketones (2) terminal alkynes from non-terminal alkynes.  
 87. (3) 1-iii, 2-i, 3-ii, 4-iii




Hence,  $H_2S_2O_8$  has O—O (peroxide) linkage.



A zwitter ion is formed by transfer of a proton from a  $-COOH$  groups to an  $-NH_2$  group.

90. (3) Combustion reactions are always exothermic as they release heat so  $\Delta H = -ve$ .  
 91. (1) a-iv, b-iii, c-ii, d-i  
 92. (4) Number of milli Equivalents of  $KMnO_4 = 0.04 \times 5 \times 50 = 10$   
 Number of milli Equivalents of  $H_2C_2O_4 = 50 \times 2 \times 0.1 = 10$   
 93. (3) Acidic character :  $HOCl < HClO_2 < HClO_3 < HClO_4$   
 Oxidising power :  $HOCl > HClO_2 > HClO_3 > HClO_4$   
 Thermal stability :  $HOCl < HClO_2 < HClO_3 < HClO_4$   
 'Cl—O' bond order :  $HOCl < HClO_2 < HClO_3 < HClO_4$   
 94. (3) Vitamin                      Deficiency disease  
 A. Vitamin- $B_{12}$                       1. Pernicious anaemia  
 B. Vitamin- $B_6$                       2. Skin disease  
 C. Vitamin-E                      3. Sterility  
 D. Vitamin-K                      4. Haemorrhagic condition  
 95. (1) a-iv b-i, c-ii, d-iii



97. (1)  $\ln \frac{K_2}{K_1} = \frac{\Delta H}{R} = \frac{1}{T_1} - \frac{1}{T_2}$

As  $K_{eq}$  is increasing with decrease in temperature, reaction is Exothermic.

98. (3) Helium provide inert atmosphere in the welding of metals of alloys that are easily oxidised. Argon is used in gas filled electric lamps, i.e., cryogenics  
 Neon is used in electric sign, i.e., advertising sign.  
 $P_4O_{10}$  is used as a valuable drying and dehydrating agent.  
 $PCl_5$  is used in organic reaction for the replacement of hydroxyl group by chlorine atom.  
 99. (4) All the above  
 100. (1)  $FeSO_4 \cdot NO$